



Idempotents and structures of rings

P.N. Ánh, G.F. Birkenmeier & L. van Wyk

To cite this article: P.N. Ánh, G.F. Birkenmeier & L. van Wyk (2016) Idempotents and structures of rings, *Linear and Multilinear Algebra*, 64:10, 2002-2029, DOI: 10.1080/03081087.2015.1134429

To link to this article: <http://dx.doi.org/10.1080/03081087.2015.1134429>



Published online: 20 Jan 2016.



Submit your article to this journal [↗](#)



Article views: 43



View related articles [↗](#)



View Crossmark data [↗](#)

Idempotents and structures of rings

P.N. Ánh^a, G.F. Birkenmeier^{b*} and L. van Wyk^c

^aRényi Institute of Mathematics, Hungarian Academy of Sciences, Budapest, Hungary;

^bDepartment of Mathematics, University of Louisiana at Lafayette, Lafayette, LA, USA;

^cDepartment of Mathematical Sciences, Stellenbosch University, Stellenbosch, South Africa

Communicated by M. Chebotar

(Received 13 September 2015; accepted 15 December 2015)

Recall that an n -by- n generalized matrix ring is defined in terms of sets of rings $\{R_i\}_{i=1}^n$, (R_i, R_j) -bimodules $\{M_{ij}\}$ and bimodule homomorphisms $\theta_{ijk} : M_{ij} \otimes_{R_j} M_{jk} \rightarrow M_{ik}$, where the set of diagonal matrix units $\{E_{ii}\}$ form a complete set of orthogonal idempotents. Moreover, an arbitrary ring with a complete set of orthogonal idempotents $\{e_i\}_{i=1}^n$ has a Peirce decomposition which can be arranged into an n -by- n generalized matrix ring R^π which is isomorphic to R . In this paper, we focus on the subclass \mathcal{T}_n of n -by- n generalized matrix rings with $\theta_{iji} = 0$ for $i \neq j$. \mathcal{T}_n contains all upper and all lower generalized triangular matrix rings. The triviality of the bimodule homomorphisms motivates the introduction of three new types of idempotents called the inner Peirce, outer Peirce and Peirce trivial idempotents. These idempotents are our main tools and are used to characterize \mathcal{T}_n and define a new class of rings called the n -Peirce rings. If R is an n -Peirce ring, then there is a certain complete set of orthogonal idempotents $\{e_i\}_{i=1}^n$ such that $R^\pi \in \mathcal{T}_n$. We show that every n -by- n generalized matrix ring R contains a subring S which is maximal with respect to being in \mathcal{T}_n and S is essential in R as an (S, S) -bisubmodule of R . This allows for a useful transfer of information between R and S . Also, we show that any ring is either an n -Peirce ring or for each $k > 1$ there is a complete set of orthogonal idempotents $\{e_i\}_{i=1}^k$ such that $R^\pi \in \mathcal{T}_k$. Examples are provided to illustrate and delimit our results.

Keywords: idempotent; generalized matrix ring; formal matrix ring; Morita Context; Peirce trivial; annihilator; bimodule; essential; ideal extending

AMS Subject Classifications: 16S50; 15A33; 16D20; 16D70

1. Introduction

Throughout this paper all rings are associative with a unity and modules are unital unless explicitly indicated otherwise.

Given a complete set of orthogonal idempotents, $\{e_i\}_{i=1}^n$, of a ring R , we can form a group direct sum,

$$R = e_1 R e_1 \oplus \cdots \oplus e_1 R e_n \oplus e_2 R e_1 \oplus \cdots \oplus e_2 R e_n \oplus \cdots \oplus e_n R e_1 \oplus \cdots \oplus e_n R e_n,$$

*Corresponding author. Email: gfb1127@louisiana.edu

called the Peirce decomposition of R . This decomposition can be arranged into an n -by- n square array, called R^π , with

$$R^\pi = \begin{bmatrix} e_1 R e_1 & e_1 R e_2 & \cdots & e_1 R e_n \\ e_2 R e_1 & e_2 R e_2 & \ddots & \vdots \\ \vdots & \ddots & \ddots & e_{n-1} R e_n \\ e_n R e_1 & \cdots & e_n R e_{n-1} & e_n R e_n \end{bmatrix}.$$

The array R^π forms a ring, where addition is componentwise and multiplication is the usual row-column matrix multiplication. Moreover, there is a ring isomorphism $h : R \rightarrow R^\pi$ defined by $h(x) = [e_i x e_j]$ for all $x \in R$. Observe that the $e_i R e_i$ are rings with unity and the $e_i R e_j$ are $(e_i R e_i, e_j R e_j)$ -bimodules. Note that the bimodule product $e_i R e_j \cdot e_j R e_k$, arising in the row-column multiplication, may be thought of as a bimodule homomorphism $\theta_{ijk} : e_i R e_j \otimes_{e_j R e_j} e_j R e_k \rightarrow e_i R e_k$ determined by the multiplication of R .

The above discussion motivates the following well-known definition:

an n -by- n generalized (or formal) matrix ring R is a square array

$$R = \begin{bmatrix} R_1 & M_{12} & \cdots & M_{1n} \\ M_{21} & R_2 & \ddots & \vdots \\ \vdots & \ddots & \ddots & M_{n-1,n} \\ M_{n1} & \cdots & M_{n,n-1} & R_n \end{bmatrix}$$

where each R_i is a ring, each M_{ij} is an (R_i, R_j) -bimodule and there exist (R_i, R_k) -bimodule homomorphisms $\theta_{ijk} : M_{ij} \otimes_{R_j} M_{jk} \rightarrow M_{ik}$ for all $i, j, k = 1, \dots, n$ (with $M_{ii} = R_i$). For $m_{ij} \in M_{ij}$ and $m_{jk} \in M_{jk}$, $m_{ij} m_{jk}$ denotes $\theta_{ijk}(m_{ij} \otimes m_{jk})$. The homomorphisms θ_{ijk} must satisfy the associativity relation: $(m_{ij} m_{jk}) m_{k\ell} = m_{ij} (m_{jk} m_{k\ell})$ for all $m_{ij} \in M_{ij}$, $m_{jk} \in M_{jk}$, $m_{k\ell} \in M_{k\ell}$ and all $i, j, k, \ell = 1, \dots, n$. Observe that θ_{iij} is determined by the ring multiplication in R_i , while θ_{ijj} and θ_{jjk} are determined by the bimodule scalar multiplications. Further information on generalized matrix rings can be found in [1].

With these conditions, addition on R is componentwise and multiplication on R is row-column matrix multiplication. A Morita context is a 2-by-2 generalized matrix ring. An n -by- n generalized upper (lower) triangular matrix ring is a generalized matrix ring with $M_{ij} = 0$ for $j < i$ ($M_{ij} = 0$ for $i < j$). Note that $\{E_{ii} \in R \mid E_{ii}$ is the matrix with $1 \in R_i$ in the (i, i) -position and 0 elsewhere, $i = 1, \dots, n\}$ is a complete set $\{E_{ii}\}_{i=1}^n$ of orthogonal idempotents in the above constructed generalized matrix ring R .

The foregoing observations allow us to consider a generalized matrix ring in two ways:

- (1) given a ring R and a complete set of orthogonal idempotents, $\{e_i\}_{i=1}^n$, then R^π is an ‘internal’ representation of R as a generalized matrix ring in terms of substructures of R ; whereas
- (2) given collections $\{R_i\}$, $\{M_{ij}\}$, and $\{\theta_{ijk}\}$, we construct a new ring from these ‘external’ components via the generalized matrix ring notion.

An important problem in the study of generalized matrix rings is: given a collection of rings $\{R_i \mid i = 1, \dots, n\}$ and bimodules $\{M_{ij} \mid i, j = 1, \dots, n, i \neq j, \text{ and each } M_{ij} \text{ is an } (R_i, R_j) \text{ - bimodule}\}$ determine the θ_{iji} ($i \neq j$) and the θ_{ijk} (i, j, k distinct) to produce an

n -by- n generalized matrix ring. We can simplify this problem by trivializing the θ_{ijk} in the following three ways (note that for $n = 2$, all three conditions coincide):

- (I) Define $\theta_{iji} = 0$, for all $i \neq j$.
- (II) Define $\theta_{ijk} = 0$, for all i, j, k pairwise distinct.
- (III) Define $\theta_{ijk} = 0$, for all $i \neq j$ and $j \neq k$ (I and II combined).

Two questions immediately arise:

- (A) Are there significant examples of generalized matrix rings with trivialized θ_{ijk} ?
- (B) How can the theory of generalized matrix rings with trivialized θ_{ijk} be used to gain insight into the theory of arbitrary generalized matrix rings?

In this paper, we consider the class of n -by- n ($n > 1$) generalized matrix rings satisfying condition (I) (i.e. $\theta_{iji} = 0$, for all $i \neq j$).

We denote this class of rings by \mathcal{T}_n .

For each generalized matrix ring R ,

we use \bar{R} to denote the ring in \mathcal{T}_n which has the same corresponding $R_i, M_{ij}, \theta_{ijk}$ as R , except that for all $i \neq j$ the homomorphisms θ_{iji} are taken to be 0 in \bar{R} .

Thus R and \bar{R} are the same ring if and only if $R \in \mathcal{T}_n$. Note that the classes of n -by- n generalized upper and lower triangular matrix rings form significant proper subclasses of \mathcal{T}_n (see Question A). Further examples are provided throughout this paper.

Observe that the triviality of the θ_{iji} motivates three new types of idempotents which appear in the internal (Peirce decomposition) generalized matrix ring representation of a ring in \mathcal{T}_2 . For $e = e^2 \in R$,

- (1) e is *inner Peirce trivial* if $eR(1 - e)Re = 0$;
- (2) e is *outer Peirce trivial* if $(1 - e)ReR(1 - e) = 0$;
- (3) e is *Peirce trivial* if e is both inner and outer trivial.

In [2] Peirce introduced the concept of an idempotent, and so we are naming certain idempotents and rings in this paper in his honour. These idempotents provide the main tools in our investigations; in particular, they are used to characterize the class \mathcal{T}_n and the class of n -Peirce rings.

In Section 3, we develop the basic properties of the inner (outer) Peirce trivial idempotents. Moreover we show that if R is a subring of a ring T and S is the subring of T generated by R and a subset \mathcal{E} of inner or outer Peirce trivial idempotents of T , then there is a useful transfer of information between R and S , e.g. R is strongly π -regular or has classical Krull dimension n if and only if so does S (Theorems 3.13, 3.14 and 3.16).

In Section 4, we begin by showing that, for a ring R with a complete set $\{e_i\}_{i=1}^n$ of orthogonal idempotents, $R^\pi \in \mathcal{T}_n$ if and only if each e_i is inner Peirce trivial (Theorem 4.2).

Next, let R be a generalized n -by- n matrix ring and take

$$\mathcal{D}(R) = \{[r_{ij}] \in R \mid r_{ij} = 0 \text{ for all } i \neq j\}$$

and

$$\mathcal{D}(R)^- = \{[r_{ij}] \in R \mid r_{ii} = 0 \text{ for all } i = 1, \dots, n\}.$$

We obtain that if $R \in \mathcal{T}_n$ then $\mathcal{D}(R)^- \trianglelefteq R$ such that $(\mathcal{D}(R)^-)^n = 0$ and $\mathcal{D}(R)$ is ring isomorphic to $R/(\mathcal{D}(R)^-)$ (Proposition 4.4).

The transfer of various ring properties (e.g. semilocal, bounded index, having a polynomial identity) between R and $\mathcal{D}(R)$ is considered when $R \in \mathcal{T}_n$. In Theorem 4.12 (one of the main results of the paper) we show that every n -by- n generalized matrix ring has subrings S maximal with respect to being in \mathcal{T}_n such that S is essential in R as an (S, S) -bimodule. This fact allows for a two-step transfer of information from $\mathcal{D}(R)$ to S (Theorem 3.16) and from S to R (Theorem 4.12 and Corollary 4.14).

In the remainder of this section, we introduce the notion of an ideal extending ring and use Theorem 4.12 and its consequences to show how this notion passes from a ring A to certain generalized matrix rings which are overrings of the n -by- n upper triangular matrix ring over A . Thus Theorem 4.12 and its corollaries provide answers to Question B.

The n -Peirce rings are introduced and investigated in Section 5. A ring R is a

1-Peirce ring

if 0 and 1 are the only Peirce trivial idempotents in R . Inductively, for a natural number $n > 1$, we say a ring R is an

n -Peirce ring

if there is a Peirce trivial idempotent e such that eRe is an m -Peirce ring for some $1 \leq m < n$ and $(1 - e)R(1 - e)$ is an $(n - m)$ -Peirce ring.

In Theorem 5.7, we show that an n -Peirce generalized matrix ring is in \mathcal{T}_n ($n > 1$) and that if R has a complete set $\{e_i\}_{i=1}^n$ of orthogonal idempotents such that each e_iRe_i is a 1-Peirce ring, then R is a k -Peirce ring for some $1 \leq k \leq n$. Example 5.2 shows that the class of n -Peirce rings is a proper subclass of \mathcal{T}_n for $n > 1$, and that any n -by- n generalized upper triangular matrix ring with prime diagonal rings is an n -Peirce ring. The class of n -Peirce rings has an advantage over \mathcal{T}_n in that for $n > 1$, an n -Peirce ring has a complete set $\{e_i\}_{i=1}^n$ of orthogonal idempotents such that each e_iRe_i is a 1-Peirce ring. In Theorem 5.11, it is also shown that if R has DCC on $\{ReR \mid e \text{ is a Peirce trivial idempotent}\}$, then R is an n -Peirce ring for some n .

As indicated in Definition 3.1, the inner (outer) Peirce trivial idempotents can be defined in a ring without a unity. Hence, many of the results in this paper can be modified to hold in rings without a unity.

2. Notation and terminology

- (1) R is Abelian – means every idempotent is central.
- (2) $\mathcal{B}(R)$, $\mathcal{P}(R)$ and $\mathcal{J}(R)$ denote the central idempotents of R , the prime radical of R and the Jacobson radical of R , respectively.
- (3) $\mathcal{S}_\ell(R) = \{e = e^2 \in R \mid Re = eRe\}$, $\mathcal{S}_r(R) = \{e = e^2 \in R \mid eR = eRe\}$.
- (4) $\text{Cen}(R)$ is the center of R .
- (5) $U(R)$ is the group of units of R .
- (6) $\langle - \rangle_R$ is the subring of R generated by $-$, and $(-)_R$ is the ideal of R generated by $-$.
- (7) $X \trianglelefteq R$ means X is an ideal of R .

- (8) $r_A(B)$ and $\ell_A(B)$ denote the right and left annihilator of B in A , respectively.
- (9) \mathbb{Z} and \mathbb{Z}_n denote the ring of integers and the ring of integers modulo n , respectively.
- (10) \mathbb{Z}^+ means the positive integers.

3. Basic properties of Peirce trivial idempotents

Definition 3.1 Let R be a ring, not necessarily with a unity, and let $e = e^2 \in R$. We say e is *inner Peirce trivial* (respectively, *outer Peirce trivial*) if $exye = exeye$ (respectively, $xe y + exeye = xeye + exey$) for all $x, y \in R$. If e is both inner and outer Peirce trivial, we say e is *Peirce trivial*.

For a ring R with a unity, e is inner (respectively, outer) Peirce trivial if and only if $eR(1 - e)Re = \{0\}$ (respectively, $(1 - e)ReR(1 - e) = \{0\}$); moreover, e is inner Peirce trivial if and only if $f = 1 - e$ is outer Peirce trivial. Let

$$\mathfrak{P}_{it}(R), \mathfrak{P}_{ot}(R) \text{ and } \mathfrak{P}_t(R)$$

denote the set of all inner Peirce trivial idempotents, all outer Peirce trivial idempotents and all Peirce trivial idempotents of R , respectively. Note that $\mathcal{B}(R) \subseteq \mathfrak{P}_t(R)$.

Example 3.2 Inner and outer Peirce trivialities are independent properties of idempotents. Let $R_1 = \mathbb{Z}, R_2 = \mathbb{Z}/8\mathbb{Z} = \mathbb{Z}_8, M_{12} = \mathbb{Z}_4, M_{21} = \mathbb{Z}_2$, together with tensor products $M_{12} \otimes_{R_2} M_{21} \cong \mathbb{Z}_2 \mapsto 0 \in R_1$ and $M_{21} \otimes_{R_1} M_{12} \cong \mathbb{Z}_2 \cong 4R_2$, respectively. Then in $R = \begin{bmatrix} R_1 & M_{12} \\ M_{21} & R_2 \end{bmatrix}$ the elements $e = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ and $f = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$ are idempotents. Moreover, e is inner Peirce trivial, but not outer Peirce trivial, and f is outer Peirce trivial, but not inner Peirce trivial.

PROPOSITION 3.3 Let $R = \begin{bmatrix} R_1 & M_{12} \\ M_{21} & R_2 \end{bmatrix} \in \mathcal{T}_2$ and assume $\alpha = \begin{bmatrix} e & m \\ n & f \end{bmatrix} \in R$.

- (1) Then $\alpha = \alpha^2$ if and only if $e = e^2, f = f^2, em + mf = m$ and $ne + fn = n$.
- (2) If $\alpha = \alpha^2$ and e and f are central idempotents, then $\alpha \in \mathfrak{P}_t(R)$. In particular, if R_1 and R_2 are commutative, then $\mathfrak{P}_t(R) = \{\alpha \in R \mid \alpha = \alpha^2\}$.

Proof The proof is a straightforward calculation using Definition 3.1. □

Note that Proposition 3.3(2) is, in general, not true when $R \in \mathcal{T}_n$ for $n > 2$ (see Example 3.9).

As a consequence of Definition 3.1, one has the following descriptions:

LEMMA 3.4 For $e^2 = e \in R$ the following claims are equivalent:

- (1) e is inner Peirce trivial.
- (2) $e\ell_R(e) = eR(1 - e)$ is a right ideal of R .
- (3) $r_R(e)e = (1 - e)Re$ is a left ideal of R .
- (4) $efge = efge$ for all idempotents $f, g \in R$.
- (5) $h : R \rightarrow eRe$, defined by $h(x) = exe$, is a surjective ring homomorphism.

- (6) $eRtRe = 0$ for all $t \in R$ such that $ete = 0$.
- (7) $ReR \subseteq \underline{\ell}_R((1 - e)Re)$.

Proof We show implication (4) \Rightarrow (1); the remaining implications are routine. For any $x, y \in R$ simple computation shows $f := e - ex + exe = ef = f^2$ and $g := e - ye + eye = ge = g^2$, whence one has by assumption $e + exye - exeye = efge = efge = e$, implying $exye = exeye$. Therefore e is inner Peirce trivial. \square

LEMMA 3.5 For $e^2 = e \in R$ the following claims are equivalent:

- (1) e is outer Peirce trivial.
- (2) $e\underline{\ell}_R(e) = eR(1 - e)$ is a left ideal of R .
- (3) $\underline{r}_R(e)e = (1 - e)Re$ is a right ideal of R .
- (4) $feg + efeg = fege + efeg$ for all idempotents $f, g \in R$.

Proof Again, we show the implication (4) \Rightarrow (1); the remaining implications are routine. For any $x, y \in R$ simple computation shows $f := e + xe - exe = fe = f^2$ and $g := e + ey - eye = eg = g^2$, whence one has by assumption $feg + efeg = fg + e = fege + efeg = f + g$. Therefore we have the equality $e + (e + xe - exe)(e + ey - eye) = e + xe - exe + e + ey - eye$, from which one can obtain, after simplification, that e is outer Peirce trivial. \square

COROLLARY 3.6 For $e^2 = e \in R$ the following claims are equivalent:

- (1) e is Peirce trivial.
- (2) $e\underline{\ell}_R(e) = eR(1 - e)$ is an ideal of R .
- (3) $\underline{r}_R(e)e = (1 - e)Re$ is an ideal of R .
- (4) $e, 1 - e \in \mathfrak{P}_{it}(R)$.

From the above results, one can see that if R is semiprime, then $\mathfrak{P}_{it}(R) = \mathfrak{P}_{ot}(R) = \mathfrak{P}_t(R) = \mathcal{B}(R)$.

LEMMA 3.7 Let $e, f \in R$ such that $e = e^2$ and $f = f^2$.

- (1) $e \in \mathfrak{P}_{it}(R)$ implies $efe = (efe)^2$, $(ef)^2 = (ef)^3$ and $(fe)^2 = (fe)^3$.
- (2) $e \in \mathfrak{P}_t(R)$ implies $fef = (fef)^2$.
- (3) $e, f \in \mathfrak{P}_{it}(R)$ implies $efe, fef \in \mathfrak{P}_{it}(R)$.
- (4) If R is a generalized matrix ring and $[e_{ij}] \in \mathfrak{P}_{it}(R)$ (resp. $\mathfrak{P}_{ot}(R)$, $\mathfrak{P}_t(R)$), then $e_{ii} \in \mathfrak{P}_{it}(R_i)$ (resp. $\mathfrak{P}_{ot}(R_i)$, $\mathfrak{P}_t(R_i)$).

Proof This proof is routine. \square

LEMMA 3.8 Let $c, e \in R$ such that $c = c^2$ and $e = e^2$.

- (1) $e \in \mathfrak{P}_{it}(R)$ if and only if $\mathfrak{P}_{it}(eRe) = eRe \cap \mathfrak{P}_{it}(R)$.
- (2) $\mathfrak{P}_t(R) \cap cRc \subseteq \mathfrak{P}_t(cRc)$.
- (3) Assume $I \trianglelefteq R$. Then $\mathfrak{P}_{it}(I) = I \cap \mathfrak{P}_{it}(R)$.

Proof

- (1) Clearly, $eRe \cap \mathfrak{P}_{it}(R) \subseteq \mathfrak{P}_{it}(eRe)$. Assume $e \in \mathfrak{P}_{it}(R)$, $c \in \mathfrak{P}_{it}(eRe)$ and $x, y \in R$. Then $cxyz = c(exye)c = c((exe)(eye))c = c(exe)c(eye)c = cxyz$. Thus $eRe \cap \mathfrak{P}_{it}(R) = \mathfrak{P}_{it}(eRe)$. Conversely, assume $eRe \cap \mathfrak{P}_{it}(R) = \mathfrak{P}_{it}(eRe)$. Since $e \in \mathfrak{P}_{it}(eRe)$, then $e \in \mathfrak{P}_{it}(R)$.
- (2) The proof of this part is straightforward.
- (3) Clearly, $I \cap \mathfrak{P}_{it}(R) \subseteq \mathfrak{P}_{it}(I)$. Let $f \in \mathfrak{P}_{it}(I)$ and $x, y \in R$. Then $fxyf = f((fx)(yf))f = f(fx)f(yf)f = fxfyf$. Therefore $\mathfrak{P}_{it}(I) = I \cap \mathfrak{P}_{it}(R)$. \square

Example 3.9 In general, for $c \in \mathfrak{P}_t(R)$, $\mathfrak{P}_t(R) \cap cRc \subsetneq \mathfrak{P}_t(cRc)$. Let R be the 3-by-3 upper triangular matrix ring over a ring A with $e = E_{22} \in R$ and $c = E_{22} + E_{33}$. Then $c \in \mathfrak{P}_t(R)$ and $e \in \mathfrak{P}_t(cRc)$, but $e \notin \mathfrak{P}_{ot}(R)$. Thus, $\mathfrak{P}_t(R) \cap cRc \subsetneq \mathfrak{P}_t(cRc)$.

In [3] (also see [4,5]), it is shown that a ring R has a generalized triangular matrix form if and only if it has a set of left (or right) triangulating idempotents. Such a set is an ordered complete set of orthogonal idempotents which are constructed from $\mathcal{S}_\ell(R)$ and $\mathcal{S}_r(R)$.

Our next result and results from Sections 4 and 5 show that $\mathfrak{P}_{it}(R)$ and $\mathfrak{P}_t(R)$ can be used to naturally extend the notion of a generalized triangular matrix ring. Moreover, the inherent symmetry in the definitions of $\mathfrak{P}_{it}(R)$ and $\mathfrak{P}_t(R)$ frees us from the ‘ordered’ condition on sets of idempotents when characterizing these natural extensions.

PROPOSITION 3.10

- (1) $\mathcal{S}_l(R) \cup \mathcal{S}_r(R) \subseteq \mathfrak{P}_t(R)$.
- (2) Let $\{e_1, \dots, e_n\}$ be a set of left or right triangulating idempotents of R . Then $\{e_1, \dots, e_n\} \subseteq \mathfrak{P}_{it}(R)$.

Proof

- (1) This part is immediate from the definitions.
- (2) From [3, p. 560 and Corollary 1.6] and Lemma 1.8, each $e_i \in \mathfrak{P}_{it}(R)$. \square

From Proposition 3.10, $\mathcal{B}(R) \subseteq \mathcal{S}_l(R) \cup \mathcal{S}_r(R) \subseteq \mathfrak{P}_t(R) \subseteq \mathfrak{P}_{it}(R) (\mathfrak{P}_{ot}(R))$. If R is semiprime these containment relations become equalities by Lemmas 3.4 and 3.5.

Example 3.11 Let A be a ring whose only idempotents are 0 and 1. Assume $0 \neq X, Y \trianglelefteq A$. Using Proposition 3.3 we obtain:

- (1) Let $R = \begin{bmatrix} A & X \\ 0 & A \end{bmatrix}$. Then $\mathcal{S}_l(R) \cup \mathcal{S}_r(R) = \mathfrak{P}_t(R) = \{e \mid e = e^2 \in R\}$.
- (2) Let $R = \begin{bmatrix} A & X \\ Y & A \end{bmatrix}$ and $XY = 0 = YX$. Then $\mathcal{S}_r(R) = \mathcal{S}_l(R) = \{0, 1\} \subsetneq \mathfrak{P}_t(R) = \{e \mid e = e^2 \in R\}$.

- (3) Let B be a subring of A with $X \trianglelefteq B$ and $R = \begin{bmatrix} A & X \\ A/X & B \end{bmatrix}$. Then $\mathcal{S}_r(R) = \mathcal{S}_l(R) = \{0, 1\} \subsetneq \mathfrak{P}_t(R) = \{e \mid e = e^2 \in R\}$.

We conclude Section 3 by showing in the next results (3.12–3.16) that given a base ring R , an overring T , and a set \mathcal{E} contained in $\mathfrak{P}_{it}(T) \cup \mathfrak{P}_{ot}(T)$, there is a significant transfer of information between R and S , where S is the subring of T generated by R and \mathcal{E} . These results indicate the importance of the inner and outer Peirce trivial idempotents.

Let S be an overring of R . We consider the following properties between prime ideals of R and S (see [6, pp. 295–296] or [7, p.292]).

- (1) *Lying over (LO)*. For any prime ideal P of R , there exists a prime ideal Q of S such that $P = Q \cap R$.
- (2) *Going up (GO)*. Given prime ideals $P_1 \subseteq P_2$ of R and Q_1 of S with $P_1 = Q_1 \cap R$, there exists a prime ideal Q_2 of S with $Q_1 \subseteq Q_2$ and $P_2 = Q_2 \cap R$.
- (3) *Incomparable (INC)*. Two different prime ideals of S with the same contraction in R are not comparable.

LEMMA 3.12 Let T be a ring, R a subring of T ,

$$\mathcal{E}_{\mathcal{P}} = \{e = e^2 \in T \mid e + \mathcal{P}(T) \text{ is central in } T/\mathcal{P}(T)\},$$

and $S = \langle R \cup \mathcal{E} \rangle_T$, where $\emptyset \neq \mathcal{E} \subseteq \mathcal{E}_{\mathcal{P}}$. Then:

- (1) $\mathfrak{P}_{it}(T) \cup \mathfrak{P}_{ot}(T) \subseteq \mathcal{E}_{\mathcal{P}}$.
- (2) If K is a prime ideal of S , then $R/(K \cap R) \cong S/K$.
- (3) *LO, GU and INC hold between R and S .*

Proof

- (1) This part follows from Lemmas 3.4 and 3.5.
- (2) and (3). The proof of these parts is similar to that in [8, Lemma 2.1] or [6, Lemma 8.3.26]. □

Recall that a ring R is *strongly π -regular* if for each x there is a positive integer n (depending on x) such that $x^n \in x^{n+1}R$.

THEOREM 3.13 Let \mathcal{C} be a property of rings such that a ring A has property \mathcal{C} if and only if every prime factor of A has property \mathcal{C} . Assume T is a ring, R is a subring of T and $S := \langle R \cup \mathcal{E} \rangle_T$, where $\emptyset \neq \mathcal{E} \subseteq \mathcal{E}_{\mathcal{P}}$, with $\mathcal{E}_{\mathcal{P}}$ as in Lemma 3.12. Then R has property \mathcal{C} if and only if S has property \mathcal{C} . In particular, R is strongly π -regular if and only if S is strongly π -regular.

Proof Assume R has property \mathcal{C} and K is a prime ideal of S . From Lemma 3.12(2), $R/(K \cap R)$ is a prime ring, and so $R/(K \cap R)$ has property \mathcal{C} . Hence S/K has property \mathcal{C} . Therefore S has property \mathcal{C} .

Conversely, assume S has property \mathcal{C} and P is a prime ideal of R . From Lemma 3.12(3), LO holds between R and S . So there exists a prime ideal Q of S such that $Q \cap R = P$. By

Lemma 3.12(2), $R/P = R/(Q \cap R) \cong S/Q$. Hence R/P has property \mathcal{C} . Therefore R has property \mathcal{C} .

From [9], a ring A is strongly π -regular if and only if every prime factor of A is a strongly π -regular ring. \square

See [10] for the definition of a special radical. Observe that the prime, Jacobson and nil radicals are included in the collection of special radicals.

THEOREM 3.14 *Let R be a subring of a ring T , $\emptyset \neq \mathcal{E} \subseteq \mathcal{E}_{\mathcal{P}}$, and $S = \langle R \cup \mathcal{E} \rangle_T$. Then:*

- (1) $\rho(R) = \rho(S) \cap R$, where ρ is any special radical.
- (2) The classical Krull dimensions of both S and R are equal.
- (3) If S is a von Neumann regular ring, then so is R .

Proof Using Lemma 3.12, the proof is similar to [8, Theorem 2.2] or [6, Theorem 8.3.28]. \square

LEMMA 3.15 *Let $T \in \mathcal{T}_n$ ($n > 1$) and*

$$\mathcal{E}_k = \{[t_{ij}] \in T \mid t_{kj} \in M_{kj} \text{ for } j \neq k, t_{kk} = 1 \in T_k \text{ and all other entries are zero}\}.$$

Then $\cup_{k=1}^n \mathcal{E}_k \subseteq \mathfrak{P}_{\text{it}}(T)$.

Proof The proof of this result is a routine but tedious application of Definition 3.1. \square

THEOREM 3.16 *Let $T \in \mathcal{T}_n$ ($n > 1$), $R = \mathcal{D}(T)$ and $\mathcal{E} = \cup_{k=1}^n \mathcal{E}_k$ (as in Lemma 3.15). Then:*

- (1) $\langle R \cup \mathcal{E} \rangle_T = S = T$.
- (2) R has property \mathcal{C} (as in Theorem 3.13) if and only if T has property \mathcal{C} .
- (3) $\rho(R) = \rho(T) \cap R$.
- (4) The classical Krull dimension of both R and T are equal.

Proof Use Lemmas 3.12 and 3.15 and Theorems 3.13 and 3.14. \square

This result extends [11, Corollary 3.6].

4. Characterization of \mathcal{T}_n

LEMMA 4.1 *Let $\{e_1, \dots, e_n\}$ be a complete set of orthogonal idempotents of R .*

- (1) $e_i \in \mathfrak{P}_{\text{it}}(R)$ if and only if $e_i R e_j R e_i = 0$ for all $j \neq i$.
- (2) $e_j \in \mathfrak{P}_{\text{ot}}(R)$ if and only if $e_i R e_j R e_k = 0$ for all $i \neq j$ and $k \neq j$.
- (3) $\{e_1, \dots, e_n\} \subseteq \mathfrak{P}_{\text{ot}}(R)$ if and only if $\{e_1, \dots, e_n\} \subseteq \mathfrak{P}_{\text{t}}(R)$.

Proof (3) follows obviously from (1) and (2). Furthermore, (1) and (2) are immediate consequences of the definition of Peirce trivial idempotents by observing

$$0 = e_i R(1 - e_i) R e_i = e_i R \left(\sum_{j \neq i} e_j \right) R e_i = \sum_{j \neq i} e_i R e_j R e_i \Leftrightarrow \forall j \neq i \ e_i R e_j R e_i = 0,$$

and

$$0 = (1 - e_j) R e_j R (1 - e_j) = \oplus_{i \neq j \neq k} e_i R e_j R e_k \Leftrightarrow \forall i \neq j \neq k \ e_i R e_j R e_k = 0. \quad \square$$

Lemma 4.1 shows remarkably that inner and outer Peirce trivial idempotents behave quite differently when they are considered together as a complete set of idempotents although their definition seems very symmetric! Lemma 4.1 shows clearly the equivalence of the first three statements in the next result.

THEOREM 4.2 *Let $\{e_1, \dots, e_n\}$ be a complete set of orthogonal idempotent elements of R . The following conditions are equivalent:*

- (1) $R^\pi \in \mathcal{T}_n$.
- (2) $e_i R e_j R e_i = 0$, for all $i \neq j$.
- (3) $\{e_1, \dots, e_n\} \subseteq \mathfrak{P}_{\text{it}}(R)$.
- (4) $\mathcal{D}(R^\pi)^-$ is a right ideal of R^π .

Proof (3) \Leftrightarrow (4) Let $X = \mathcal{D}(R^\pi)^-$. Observe that the (i, i) -position of $X R^\pi$ is $\sum_{k \neq i} e_i R_k R e_i = e_i R(1 - e_i) R e_i$; and the (i, j) -position of $X R^\pi$ is $\sum_{k \neq i, k \neq j} e_i R e_k R e_j \subseteq e_i R e_j$. Therefore $X R^\pi \subseteq X$ if and only if each $e_i \in \mathfrak{P}_{\text{it}}(R)$. \square

Observe that from Lemma 3.4 and Theorem 4.2, any property that is preserved by a surjective ring homomorphism passes from a ring in \mathcal{T}_n to its diagonal rings.

COROLLARY 4.3 *Let R be an n -by- n generalized matrix ring. Then the following conditions are equivalent:*

- (1) $R \in \mathcal{T}_n$.
- (2) Let $[a_{ij}], [b_{ij}] \in R$ with $[c_{ij}] = [a_{ij}][b_{ij}]$. Then $c_{ii} = a_{ii}b_{ii}$ for all i and j .
- (3) $\{E_{ii} \in R \mid i = 1, \dots, n\} \subseteq \mathfrak{P}_{\text{it}}(R)$.

Thus \mathcal{T}_n is exactly the class of n -by- n generalized matrix rings in which the diagonal entries of the product of two matrices is completely determined by the corresponding entries of the diagonals of the factor matrices.

Note that the idempotents in a generalized matrix ring are not characterized. However, for $R \in \mathcal{T}_n, e = e^2 \in R$ if and only if $e = [e_{ij}]$, where $e_{ii} = e_{ii}^2$ and $e_{ij} = \sum_{k=1}^n e_{ik}e_{kj}$ for $i \neq j$.

PROPOSITION 4.4 *Assume $\{e_1, \dots, e_n\}$ is a complete set of orthogonal idempotents of R , and $X = \mathcal{D}(R^\pi)^-$.*

- (1) *If $\{e_1, \dots, e_n\} \subseteq \mathfrak{P}_{\text{it}}(R)$, then $X^n = 0$ and $\oplus_{i=1}^n R_i$ is a homomorphic image of R^π with kernel X , where $R_i = e_i R e_i$.*
- (2) *If $\{e_1, \dots, e_n\} \subseteq \mathfrak{P}_t(R)$, then $X^2 = 0$.*

Proof

- (1) Observe that the (i, j) -position of X^{n-1} is a sum of terms where each term is an element of $e_i Re_{\alpha_1} Re_{\alpha_2} R \cdots e_{\alpha_{n-2}} Re_j$ where $\alpha_k \in \{1, \dots, n\} - \{i, j\}$. Since $\{e_1, \dots, e_n\} \subseteq \mathfrak{P}_{\text{it}}(R)$, it follows that $X^n = 0$. The second part follows from Lemma 3.4(5) and Theorem 4.2.
- (2) This part follows from Lemma 4.1(2). □

For more details on nilpotent elements and ideals in this context, see [12].

Example 4.5 Let A be a ring and $X, Y \trianglelefteq A$ such that $X^2 \subseteq Y$. Take

$$R = \begin{bmatrix} A & X & Y \\ X & A & X \\ Y & X & A \end{bmatrix}.$$

Then routine calculation yields:

- (1) $E_{22} \in \mathfrak{P}_{\text{t}}(R)$ if and only if $X^2 = 0$.
- (2) $\{E_{11}, E_{22}, E_{33}\} \subseteq \mathfrak{P}_{\text{it}}(R)$ if and only if $X^2 = 0 = Y^2$.
- (3) $\{E_{11}, E_{22}, E_{33}\} \subseteq \mathfrak{P}_{\text{t}}(R)$ if and only if $X^2 = 0 = Y^2$ and $XY = 0 = YX$.

For an illustration of (2) and (3), let B be a ring and $A = B[x, y]/(x^2, y^2)$ and $A = B[x, y]/(x^2, y^2, xy)$, respectively.

The next three results (4.6–4.8) indicate a transfer of important ring properties between $\mathcal{D}(R)$ and $R \in \mathcal{T}_n$.

COROLLARY 4.6 *Let $R \in \mathcal{T}_n$ ($n > 1$). Then $\mathcal{D}(R)$ satisfies each of the following conditions if and only if R does so:*

- (1) *semilocal,*
- (2) *semiperfect,*
- (3) *left (or right) perfect,*
- (4) *semiprimary,*
- (5) *bounded index (of nilpotence).*

Proof Observe that if R satisfies any of (1)–(5), then so does eRe for each $e = e^2 \in R$. Hence, if R satisfies any of (1)–(5), then so does $\mathcal{D}(R)$.

Conversely, first assume that $\mathcal{D}(R)$ is semilocal. From Proposition 4.4(1) we have that $\mathcal{D}(R)^- \trianglelefteq R$, $R/(\mathcal{D}(R)^-) \cong \mathcal{D}(R)$ and $\mathcal{D}(R)^- \subseteq \mathcal{J}(R)$. Therefore, $R/\mathcal{J}(R) \cong (R/(\mathcal{D}(R)^-))/(\mathcal{J}(R)/(\mathcal{D}(R)^-)) \cong \mathcal{D}(R)/\mathcal{J}(\mathcal{D}(R))$ is semisimple artinian. Thus R is semilocal. Parts (2)–(4) are proved similarly. Now assume $\mathcal{D}(R)$ has bounded index k , and let v be a nilpotent element of R . Then $v = d + x$ where $d \in \mathcal{D}(R)$ and $x \in \mathcal{D}(R)^-$ and $v^m = 0$. Then $0 = v^m = d^m + y$ where $y \in \mathcal{D}(R)^-$. So $d^m = -y \in \mathcal{D}(R)^-$. Hence d is nilpotent, so $d^k = 0$. Then $v^k = d^k + w = w \in \mathcal{D}(R)^-$. Hence $v^{kn} = 0$. Thus R has bounded index less than or equal to kn . □

Corollary 4.6(3) extends [11, Corollary 3.8]. In [13], the authors determine several generalizations of the condition that a ring satisfies a polynomial identity. With these generalizations they were able to extend classical theorems by Armendariz and Steenberg, Fisher, Kaplansky, Martindale, Posner and Rowen. Two of these generalizations are: (1) a ring R is an *almost PI-ring* if every prime factor ring of R is a PI-ring; (2) R is an *intrinsically PI-ring* if every nonzero ideal contains a nonzero PI-ideal of R .

COROLLARY 4.7 Let $R \in \mathcal{T}_n$ ($n > 1$). Then:

- (1) $\mathcal{D}(R)$ satisfies a PI if and only if R does so.
- (2) If $\mathcal{D}(R)$ is commutative, then R satisfies $(xy - yx)^n = 0$ for all $x, y \in R$.
- (3) $\mathcal{D}(R)$ is almost PI if and only if R is almost PI.
- (4) If $\mathcal{D}(R)$ is intrinsically PI, then R is intrinsically PI.

Proof

- (1) Since subrings of a PI-ring are PI-rings, if R is a PI-ring then so is $\mathcal{D}(R)$. Conversely, assume $\mathcal{D}(R)$ is a PI-ring which satisfies the PI p . Then R satisfies p^n .
- (2) This part follows from (1).
- (3) and (4). By Proposition 4.4 we have $\mathcal{D}(R)^- \trianglelefteq R$, $R/(\mathcal{D}(R)^-) \cong \mathcal{D}(R)$ and $(\mathcal{D}(R)^-)^n = 0$. Now (3) follows from [13, Theorem 1.6(i)], and (4) follows from [13, Theorem 1.6(ii)]. □

Let R be an n -by- n generalized matrix ring, and let

$$\text{UT}(R) \text{ and } \text{LT}(R)$$

be the n -by- n upper and lower generalized triangular matrix rings, respectively, formed from R . Our next result shows that elements of \mathcal{T}_n are subdirect products of generalized triangular matrix rings.

PROPOSITION 4.8 Let $R \in \mathcal{T}_n$ ($n > 1$). Then there is a ring monomorphism $\psi : R \rightarrow \text{UT}(R) \times \text{LT}(R)$ such that R is a subdirect product of $\text{UT}(R)$ and $\text{LT}(R)$.

Proof Let $[m_{ij}] \in R$. Define $\psi([m_{ij}]) = ([a_{ij}], [b_{ij}])$ where

$$a_{ij} = \begin{cases} m_{ij}, & \text{for } j \geq i \\ 0, & \text{elsewhere,} \end{cases} \quad \text{and} \\ b_{ij} = \begin{cases} m_{ij}, & \text{for } i \geq j \\ 0, & \text{elsewhere.} \end{cases}$$

A routine argument yields that ψ is a ring monomorphism and that R is a subdirect product of $\text{UT}(R)$ and $\text{LT}(R)$. □

Definition 4.9 Let R be an n -by- n generalized matrix ring. Let R^{la} denote the lower annihilating subring

$$\left[\begin{array}{cccc} R_1 & M_{12} & \cdots & M_{1n} \\ \underline{r}_{M_{21}}(M_{12}) \cap \underline{\ell}_{M_{21}}(M_{12}) & R_2 & \ddots & \vdots \\ \vdots & \ddots & \ddots & M_{n-1,n} \\ \underline{r}_{M_{n1}}(M_{1n}) \cap \underline{\ell}_{M_{n1}}(M_{1n}) & \cdots & \underline{r}_{M_{n,n-1}}(M_{n-1,n}) \cap \underline{\ell}_{M_{n,n-1}}(M_{n-1,n}) & R_n \end{array} \right]$$

of R , and let R^{ua} denote the upper annihilating subring

$$\left[\begin{array}{cccc} R_1 & \underline{r}_{M_{12}}(M_{21}) \cap \underline{\ell}_{M_{12}}(M_{21}) & \cdots & \underline{r}_{M_{1n}}(M_{n1}) \cap \underline{\ell}_{M_{1n}}(M_{n1}) \\ M_{21} & R_2 & \ddots & \vdots \\ \vdots & \ddots & \ddots & \underline{r}_{M_{n-1,n}}(M_{n,n-1}) \cap \underline{\ell}_{M_{n-1,n}}(M_{n,n-1}) \\ M_{n1} & \cdots & M_{n,n-1} & R_n \end{array} \right]$$

of R .

Note that R^{la} and R^{ua} are subrings of both R and \bar{R} . Moreover, if R is the n -by- n matrix ring over a ring A , then R^{la} and R^{ua} are the n -by- n upper and lower triangular matrix rings over A , respectively.

Example 4.10 Let A and B be rings. Let $R = \begin{bmatrix} A \times B & A \times \{0\} \\ A \times B & A \times B \end{bmatrix}$. Then $R^{la} = \begin{bmatrix} A \times B & A \times \{0\} \\ \{0\} \times B & A \times B \end{bmatrix}$, and $R^{ua} = \begin{bmatrix} A \times B & \{0\} \\ A \times B & A \times B \end{bmatrix}$.

LEMMA 4.11 Let R be an n -by- n generalized matrix ring. Then $\text{Cen}(R) = \text{Cen}(\bar{R}) = \{[c_{ij}] \in R \mid c_{ij} = 0 \text{ if } i \neq j, c_{ii} \in \text{Cen}(R_i) \text{ for all } i, \text{ and } c_{ii}m_{ij} = m_{ij}c_{jj} \text{ for all } m_{ij} \in M_{ij} \text{ if } i \neq j\}$.

Proof First show the result holds when $n = 2$. For the n -by- n case, block the matrix ring into a 2-by-2 generalized matrix ring and use induction. □

Let S be a subring of a ring R . It is well known (see [14, p. 26]) that the (S, S) -bimodule structure of S and R is equivalent to the right T -module structure of S and R , respectively, where $T = S^{\text{op}} \otimes_{\mathbb{Z}} S$, with S^{op} denoting the opposite ring of S .

The next three results (4.12–4.14) indicate the transfer of significant information between an n -by- n generalized matrix ring and certain subrings which are maximal with respect to being in \mathcal{T}_n .

In particular, Theorem 4.12 shows that for any ring R with a complete set of orthogonal idempotents $\{e_i\}_{i=1}^n$ ($n > 1$) there are subrings S containing $\{e_i\}_{i=1}^n$ which are maximal with respect to S^π being in \mathcal{T}_n and S_T is right essential in R_T , where $T = S^{\text{op}} \otimes_{\mathbb{Z}} S$. Moreover, this result and its consequences provide a connection between the structure of an arbitrary generalized matrix ring and the structure of rings in \mathcal{T}_n (see Question B in the introduction).

THEOREM 4.12 Let R be an n -by- n generalized matrix ring, and S denotes either R^{la} or R^{ua} .

- (1) S is a subring of R maximal with respect to being in \mathcal{T}_n .
- (2) Let $0 \neq y \in R$. Then either $0 \neq syE_{jj} \in S$ or $0 \neq E_{ii}yt \in S$ for some $s, t, E_{ii}, E_{jj} \in S$.
- (3) Every nonzero (S, S) -bisubmodule of R has nonzero intersection with S . Thus every nonzero ideal of R has nonzero intersection with S , and S_T is right essential in R_T where $T = S^{\text{op}} \otimes_{\mathbb{Z}} S$.
- (4) $\text{Cen}(R) = \text{Cen}(R^{\text{la}}) \cap \text{Cen}(R^{\text{ua}}) \subseteq \text{Cen}(\mathcal{D}(R))$.
- (5) $U(\bar{R}) = \{u + x \mid u \in U(\mathcal{D}(R)) \text{ and } x \in \mathcal{D}(\bar{R})^-\}$, and
 $U(S) = \{u + y \mid u \in U(\mathcal{D}(R)) \text{ and } y \in \mathcal{D}(S)^-\} \subseteq U(R)$.

Proof

- (1) Suppose that there is a subring Y of R such that S is properly contained in Y . Assume $S = R^{\text{la}}$. Then there exists $y \in Y$ with an entry y_{ij} for some i, j with $i > j$ such that $y_{ij} \notin \underline{r}_{M_{ij}}(M_{ji}) \cap \underline{\ell}_{M_{ij}}(M_{ji})$. Then either $y_{ij} \notin \underline{\ell}_{M_{ij}}(M_{ji})$, or $y_{ij} \in \underline{\ell}_{M_{ij}}(M_{ji})$ but $y_{ij} \notin \underline{r}_{M_{ij}}(M_{ji})$.

If $y_{ij} \notin \underline{\ell}_{M_{ij}}(M_{ji})$, there exists $k \in M_{ji}$ such that $y_{ijk} \neq 0$. Let t be the n -by- n matrix with k in the (j, i) -position and zero elsewhere. Then $t \in S \subseteq Y$, and so $0 \neq yt \in Y$. However, since $0 \neq y_{ijk} \in M_{ij}M_{ji} \in R_i$, we have that $Y \notin \mathcal{T}_n$.

If $y_{ij} \in \underline{\ell}_{M_{ij}}(M_{ji})$ but $y_{ij} \notin \underline{r}_{M_{ij}}(M_{ji})$, then there exists $h \in M_{ji}$ such that $hy_{ij} \neq 0$. Let s be the n -by- n matrix with h in the (j, i) -position and zero elsewhere. Then $s \in S \subseteq Y$, and so $0 \neq sy \in Y$. However, since $0 \neq hy_{ij} \in M_{ji}M_{ij} \in R_j$, it follows that $Y \notin \mathcal{T}_n$.

Therefore S is a subring of R maximal with respect to being in \mathcal{T}_n . The argument when $S = R^{\text{ua}}$ is similar.

- (2) Again let $S = R^{\text{la}}$ and $0 \neq y \in R$. If $y \in S$, we are finished. So assume $y \notin S$. Then as in part (1) there exists an entry y_{ij} of y for some i, j with $i > j$ such that $y_{ij} \notin \underline{r}_{M_{ij}}(M_{ji}) \cap \underline{\ell}_{M_{ij}}(M_{ji})$. As in part (1), we obtain $s, t \in S$. Then $0 \neq syE_{jj} \in S$ or $0 \neq E_{ii}yt \in S$. The argument when $S = R^{\text{ua}}$ is similar.
- (3) This part is a consequence of (2).
- (4) This part follows from Lemma 4.11.

- (5) By Proposition 4.4(1), $\mathcal{D}(\bar{R})^- \trianglelefteq \bar{R}$ such that $(\mathcal{D}(\bar{R})^-)^n = 0$. Let $u \in \mathcal{D}(R)$ and $x \in \mathcal{D}(\bar{R})^-$. Then $(u+x)(u^{-1}+x) = 1+ux+xu^{-1}+x^2$. But $ux+xu^{-1}+x^2 \in \mathcal{D}(\bar{R})$, hence $(u+x)(u^{-1}+x) = w$ where $w \in U(\bar{R})$. Therefore $u+x \in U(\bar{R})$. Now assume $v \in U(\bar{R})$. Then there exist $d \in \mathcal{D}(\bar{R})$ and $y \in \mathcal{D}(\bar{R})^-$ such that $v = d+y$. So $1 = dv^{-1} + yv^{-1}$. Hence $dv^{-1} = 1 - yv^{-1}$. Since $yv^{-1} \in \mathcal{D}(\bar{R})^-$, $dv^{-1} \in U(\bar{R})$. So $d \in U(\bar{R})$. Therefore $U(\bar{R}) = \{u+x \mid u \in U(\mathcal{D}(R)) \text{ and } x \in \mathcal{D}(\bar{R})^-\}$.

The remainder of the proof is due to the above argument and the fact that S is a subring of R and \bar{R} . □

Note that if $n = 2$, then in Theorem 4.12(2), there is no need for the E_{ii} and E_{jj} .

QUESTION: When is $U(R)$ generated by $U(R^{\text{la}}) \cup U(R^{\text{ua}})$?

COROLLARY 4.13 *Let R be an n -by- n generalized matrix ring, S denotes R^{la} or R^{ua} and $T = S^{\text{op}} \otimes_{\mathbb{Z}} S$. Then:*

- (1) S is maximal among subrings Y of R for which $\{E_{ii}\}_{i=1}^n \subseteq \mathfrak{P}_{\text{it}}(R)$.
- (2) The sum of the minimal ideals of S equals $\text{Soc}(S_T) = \text{Soc}(R_T)$.
- (3) The uniform dimension of ${}_S S_S$ equals the uniform dimension of ${}_S R_S$ equals the uniform dimension of S_T equals the uniform dimension of R_T .

Proof

- (1) This part is a consequence of Theorems 4.2 and 4.12.
- (2) and (3). These parts are consequences of Theorem 4.12(3). \square

Our next result demonstrates that useful information can be transferred from the diagonal rings R_i of a generalized matrix ring R to R itself via Theorems 3.16 and 4.12. Recall from [15,16] that an n -by- n ($n > 1$) matrix ring over a strongly π -regular ring is not, in general, a strongly π -regular ring.

COROLLARY 4.14 *Let R be an n -by- n generalized matrix ring, and $S = R^{\text{la}}$ or R^{ua} . If $\mathcal{D}(R)$ is strongly π -regular, then for each $0 \neq y \in R$ either:*

- (1) $y \in S$, in which case $y^n \in y^{n+1}S \subseteq y^{n+1}R$ for some positive integer n ; or
- (2) $y \notin S$, in which case either $0 \neq syE_{jj} \in S$ and $(syE_{jj})^m \in (syE_{jj})^{m+1}S \subseteq (syE_{jj})^{m+1}R$, or $0 \neq E_{ii}yv \in S$ and $(E_{ii}yv)^k \in (E_{ii}yv)^{k+1}S \subseteq (E_{ii}yv)^{k+1}R$ for some $s, v, E_{ii}, E_{jj} \in S$ and positive integers k, m, n .

Proof The proof follows from Theorems 3.16 and 4.12(2). \square

Thus if $\mathcal{D}(R)$ is strongly π -regular, then R is ‘almost’ strongly π -regular.

Next we introduce the notion of an ideal extending ring. The ideal extending condition is shown to be a Morita invariant. Moreover, it is shown that important classes of rings have this property. For example, the semiprime quasi-Baer rings are ideal extending, so this insures that every semiprime ring has a hull which is ideal extending (see Proposition 4.16). The class of semiprime quasi-Baer rings includes the local multiplier C^* -algebras which means that every C^* -algebra can be embedded into its local multiplier C^* -algebra which is an ideal extending ring.

As applications of the results in 4.12–4.14 we show in 4.16–4.21 that the ideal extending property transfers from a ring A to a certain type of overring of A which is in \mathcal{T}_n .

Let X and Y be both left or both right ideals of a ring R with $X \subseteq Y$. Then X is *ideal essential* in Y if for each $0 \neq I \trianglelefteq R$ such that $I \subseteq Y$, then $0 \neq X \cap I$. Note that if R is a semiprime ring and $X, Y \trianglelefteq R$ with $X \subseteq Y$, then X is ideal essential in Y if and only if X is right or left essential in Y .

Definition 4.15 We say R is *ideal extending* if for each $X \trianglelefteq R$ there is an $e \in \mathcal{B}(R)$ such that X is ideal essential in eR .

Note that every nonzero ideal of R is ideal essential in R if and only if $\mathcal{B}(R) = \{0, 1\}$ and R is ideal extending. Some immediate examples of ideal extending rings are: R is a prime ring, R is an Abelian (i.e. every idempotent is central) right extending ring (e.g. R

is a direct sum of commutative uniform rings (see [17]), or $R = \begin{bmatrix} A & M \\ 0 & A \end{bmatrix}$ where A is a prime ring and $M \trianglelefteq A$.

The next result shows that the class of ideal extending rings is quite extensive. See [6,8] for undefined terminology.

PROPOSITION 4.16 *Assume R is a semiprime ring. Then:*

- (1) R is ideal extending if and only if R is quasi-Baer if and only if R_R is FI-extending.
- (2) R has an ideal extending hull.

Proof

- (1) See [18, Theorem 4.7] or [6, Theorem 3.2.37].
- (2) This part follows from (1) and [8, Theorem 3.3] or [6, Theorem 8.3.17]. □

From Proposition 4.16 it follows that every von Neumann algebra and every local multiplier algebra of a C^* -algebra are ideal extending as rings (see [19] and [8, pp. 345–347] or [6, pp. 380–407]).

THEOREM 4.17

- (1) Let $\{R_i | i \in I\}$ be a set of rings. Then $\prod_{i \in I} R_i$ is ideal extending if and only if each R_i is so.
- (2) The ideal extending property is a Morita invariant.

Proof

- (1) The proof of this part is routine.
- (2) Assume that R is ideal extending. Then a straightforward argument shows that R is ideal extending if and only if the ring of n -by- n matrices over R is ideal extending. Let e be a full idempotent of R (i.e. $ReR = R$) and $0 \neq K \trianglelefteq eRe$. Then RKR is ideal essential in cR for some $c \in \mathcal{B}(R)$. Hence $K \subseteq ec(eRe)$, where $ec \in \mathcal{B}(eRe)$. Let $0 \neq X \trianglelefteq eRe$ such that $X \subseteq ec(eRe)$. Then $RXR \subseteq cR$, so $0 \neq Y = RXR \cap RKR$. If $eYe \neq 0$, there exists $y \in Y$ such that $0 \neq eye = \sum r_\alpha x_\alpha s_\alpha = \sum t_\beta k_\beta v_\beta$, where $r_\alpha, s_\alpha, t_\beta, v_\beta \in R, x_\alpha \in X$ and $k_\beta \in K$. So $x_\alpha = ex_\alpha e$ and $k_\beta = ek_\beta e$. Hence $0 \neq eye = \sum er_\alpha(ex_\alpha e)s_\alpha e = \sum et_\beta(ek_\beta e)v_\beta e \in X \cap K$.

Now assume $eYe = 0$. Since e is full, $1 = \sum a_j e b_j$ for some $a_j, b_j \in R$. Let $0 \neq w \in Y$. Then $w = 1w1 = (\sum a_j e b_j)w(\sum a_j e b_j)$. So there exists j_1, j_2 such that $a_{j_1} e b_{j_1} w a_{j_2} e b_{j_2} \neq 0$, otherwise $w = 0$, a contradiction. Hence $e b_{j_1} w a_{j_2} e \neq 0$, contrary to $eYe = 0$. Thus eRe is ideal extending. By [20, Corollary 18.35], the ideal extending property is a Morita invariant. □

PROPOSITION 4.18 *Let R be an n -by- n generalized matrix ring and $S = R^{la}$.*

- (1) If ${}_S X_S \leq {}_S R_S$ and $X \cap S$ is ideal essential in eS , for some $e \in \mathcal{S}_r(S)$, then X essential in eR as an (S, S) -bisubmodule.
- (2) If S is an ideal extending ring, then for each $X \trianglelefteq R$ there is an $e \in \mathcal{B}(S)$ such that X is ideal essential in eR .

Proof

- (1) Assume $(1 - e)X \neq 0$. Since $1 - e \in \mathcal{S}_\ell(S)$, $(1 - e)X$ is an (S, S) -bisubmodule of R . By Theorem 4.12(3), $0 \neq (1 - e)X \cap S \subseteq X \cap S \subseteq eS$, a contradiction. Then $X \subseteq eR$. Let $0 \neq {}_S Y_S \leq {}_S R_S$ such that $Y \subseteq eR$ and $Y \cap X = 0$. Hence $0 \neq Y \cap S \subseteq eS$ and $Y \cap S \trianglelefteq S$, a contradiction.
- (2) This part is a consequence of (1). □

Example 4.19

- (1) This example illustrates Proposition 4.18(1). Let $R = \begin{bmatrix} \mathbb{Z} \times \mathbb{Z}_4 & \mathbb{Z} \times 2\mathbb{Z}_4 \\ \mathbb{Z} \times \{0\} & \mathbb{Z} \times \mathbb{Z} \end{bmatrix}$, and let $S = R^{la}$. Then $S = \begin{bmatrix} \mathbb{Z} \times \mathbb{Z}_4 & \mathbb{Z} \times 2\mathbb{Z}_4 \\ \{0\} \times \{0\} & \mathbb{Z} \times \mathbb{Z} \end{bmatrix}$. Take $X = \begin{bmatrix} \{0\} \times \{0\} & \{0\} \times \{0\} \\ \{0\} \times \{0\} & \{0\} \times 2\mathbb{Z} \end{bmatrix} \trianglelefteq R$ and $e = \begin{bmatrix} (0, 0) & (0, 0) \\ (0, 0) & (0, 1) \end{bmatrix}$. Then $e \in \mathcal{S}_r(S)$ and ${}_S X_S$ is essential as an (S, S) -bisubmodule of eR . Note that $e \notin \mathcal{B}(S)$.
- (2) This example shows that in Proposition 4.18(2), S cannot be replaced by $D(R)$. Let $R = \begin{bmatrix} \mathbb{Z}_4 & 2\mathbb{Z}_4 \\ 0 & \mathbb{Z}_4 \end{bmatrix}$ (note that $R = S = R^{la}$). Then $D(R)$ is a commutative self-injective ring, hence it is ideal extending. However $\mathcal{B}(R) = \{0, 1\}$, but $\begin{bmatrix} 2\mathbb{Z}_4 & 0 \\ 0 & 0 \end{bmatrix}$ and $\begin{bmatrix} 0 & 2\mathbb{Z}_4 \\ 0 & 0 \end{bmatrix}$ are ideals of R whose intersection is zero. Therefore R is not ideal extending.

LEMMA 4.20 *If A is an ideal extending ring, then R is ideal extending where R is the n -by- n upper triangular matrix ring over A .*

Proof Let $X \trianglelefteq R$. Then

$$X = \begin{bmatrix} X_{11} & X_{12} & \cdots & X_{1n} \\ 0 & X_{22} & & X_{2n} \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \cdots & 0 & X_{nn} \end{bmatrix},$$

where each $X_{ij} \trianglelefteq A$, $X_{ii} \subseteq X_{i,i+1} \subseteq \cdots \subseteq X_{in}$ and $X_{ii} \subseteq X_{i-1,i} \subseteq \cdots \subseteq X_{1i}$ for all $1 \leq i \leq n$. Observe $e \in \mathcal{B}(R)$ if and only if $e = c1_R$, for some $c \in \mathcal{B}(A)$. There exists $f \in \mathcal{B}(A)$ such that X_{1n} is ideal essential in fA . Then a routine argument shows that X is ideal essential in $(f1_R)R$. □

The following corollary is an application of Proposition 4.18 (hence of Theorem 4.12).

COROLLARY 4.21 Let A be a ring and R the n -by- n generalized matrix ring of the form

$$R = \begin{bmatrix} A & A & \cdots & A \\ X_{21} & A & \cdots & A \\ \vdots & \ddots & \ddots & \vdots \\ X_{n1} & \cdots & X_{n,n-1} & A \end{bmatrix},$$

where $X_{ij} = A$ for $i \leq j$, $X_{ij} \trianglelefteq A$ for $j < i$, $X_{j1} \subseteq X_{j2} \subseteq \cdots \subseteq X_{jn}$ and $X_{ni} \subseteq X_{n-1,i} \subseteq \cdots \subseteq X_{1i}$, for all $1 \leq i \leq n$ and $1 \leq j \leq n$. Then A is ideal extending if and only if R is ideal extending.

Proof (\Rightarrow) Assume A is ideal extending. Observe that R^{la} is the n -by- n upper triangular matrix ring over A . By Lemma 4.20, R^{la} is ideal extending. From Proposition 4.18(2), R is ideal extending.

(\Leftarrow) Assume R is ideal extending. Let $X \trianglelefteq A$ and Y the set of n -by- n matrices over X . Then $Y \trianglelefteq R$. So there exists $e \in \mathcal{B}(R)$ such that Y is ideal essential in eR and $e = c1_R$ where $c \in \mathcal{B}(A)$. Then X is ideal essential in cA . Therefore A is ideal extending. \square

Assume R is ring isomorphic to R_m and to R_n , where R_m is an m -by- m generalized matrix ring, and R_n is an n -by- n generalized matrix ring, with $0 < m < n$. One may naturally ask:

- (1) If $R_m \in \mathcal{T}_m$, must $R_n \in \mathcal{T}_n$?
- (2) If $R_n \in \mathcal{T}_n$, must $R_m \in \mathcal{T}_m$?

The following example shows that, in general, neither question has an affirmative answer.

Example 4.22 Let A be a ring.

- (1) Let

$$R = \begin{bmatrix} A & A & A \\ A & A & A \\ 0 & 0 & A \end{bmatrix}.$$

Then $R \notin \mathcal{T}_3$, because $E_{22} \notin \mathfrak{P}_{it}(R)$ by Theorem 4.2.

Let $R_1 = \begin{bmatrix} A & A \\ A & A \end{bmatrix}$, $M_{12} = \begin{bmatrix} A \\ A \end{bmatrix}$, $M_{21} = [0 \ 0]$ and $R_2 = A$. Then $R \cong \begin{bmatrix} R_1 & M_{21} \\ M_{21} & R_2 \end{bmatrix} \in \mathcal{T}_2$.

- (2) Let

$$R = \begin{bmatrix} A & A & A & A \\ 0 & A & 0 & 0 \\ 0 & A & A & A \\ 0 & A & 0 & A \end{bmatrix}.$$

Then $R \in \mathcal{T}_4$.

$$\text{Let } R_1 = \begin{bmatrix} A & A \\ 0 & A \end{bmatrix}, M_{12} = \begin{bmatrix} A & A \\ 0 & 0 \end{bmatrix}, M_{21} = \begin{bmatrix} 0 & A \\ 0 & A \end{bmatrix} \text{ and } R_2 = \begin{bmatrix} A & A \\ 0 & A \end{bmatrix}.$$

$$\text{Then } R \cong \begin{bmatrix} R_1 & M_{12} \\ M_{21} & R_2 \end{bmatrix} \notin \mathcal{T}_2, \text{ since } M_{12}M_{21} \neq 0.$$

PROPOSITION 4.23 *Let R be a ring with a complete set $\{e_i\}_{i=1}^n$ ($n > 1$) of orthogonal idempotents. If $\{e_i\}_{i=1}^n \subseteq \mathfrak{P}_t(R)$, then any partition of R^π into an m -by- m block form is in \mathcal{T}_m where $m \leq n$.*

Proof The proof is straightforward. □

Note that in Proposition 4.23 if $n = 3$, then we can replace $\mathfrak{P}_t(R)$ with $\mathfrak{P}_{it}(R)$.

QUESTION: Let R be a ring with a complete set $\{e_i\}_{i=1}^n$ ($n > 1$) of orthogonal idempotents. What are necessary and sufficient conditions so that any partition of R^π into m -by- m block form is in \mathcal{T}_m for $1 < m \leq n$?

5. n -Peirce rings

Definition 5.1 A ring R is called a *1-Peirce ring* if $\mathfrak{P}_t(R) = \{0, 1\}$, with $0 \neq 1$. Inductively, for a natural number $n > 1$, a ring R is called an *n -Peirce ring* if there is an $e \in \mathfrak{P}_t(R)$ such that eRe is an m -Peirce ring for some $1 \leq m < n$ and $(1 - e)R(1 - e)$ is an $(n - m)$ -Peirce ring.

Example 5.2

- (1) If R_R is indecomposable or R is prime, then R is 1-Peirce. In fact, if R is semiprime then R is 1-Peirce if and only if R is indecomposable (as a ring).
- (2) If R is an n -by- n generalized upper (lower) triangular matrix ring with 1-Peirce diagonal rings, then R is a n -Peirce ring.
- (3) If R has a complete set of n orthogonal primitive idempotents which are Peirce trivial, then R is an n -Peirce ring (e.g. see Example 4.5(3)).
- (4) Example 4.5(2) is a 3-Peirce ring that has a complete set of three primitive idempotents which are inner Peirce trivial but not all of them are Peirce trivial.
- (5) Let I be an infinite index set, for each $i \in I$ let A_i be a ring with only trivial idempotents, and $A = \prod_{i \in I} A_i$. Assume $R = \begin{bmatrix} A & X \\ 0 & X \end{bmatrix}$, where X is a nonzero ideal of A . Then R is in \mathcal{T}_2 , but R is not an n -Peirce ring for any positive integer n .

From Example 5.2(5) and Theorem 5.7, we see that the class of n -Peirce n -by- n generalized matrix rings is a proper subclass of the class \mathcal{T}_n for $n > 1$. Also, due to the symmetry of Peirce idempotents (i.e. $e \in \mathfrak{P}_t(R)$ if and only if $1 - e \in \mathfrak{P}_t(R)$) the class of n -Peirce rings exhibits better behavior than \mathcal{T}_n with respect to finiteness conditions.

THEOREM 5.3 *Let R be a ring. Then either:*

- (1) R is a 1-Peirce ring;

- (2) R is an n -Peirce ring for $n > 1$, and for each $k \in \mathbb{Z}^+$ with $1 < k \leq n$ there exists a complete set of orthogonal idempotents, $\{e_i\}_{i=1}^k$, such that $R^\pi \in \mathcal{T}_k$; or
- (3) for each integer k with $k > 1$, there exists a complete set of orthogonal idempotents, $\{e_i\}_{i=1}^k$, such that $R^\pi \in \mathcal{T}_k$.

Proof Observe that $\{0, 1\} \subseteq \mathfrak{P}_t(R)$, and $\{0, 1\} = \mathfrak{P}_t(R)$ if and only if R is a 1-Peirce ring. If $\{0, 1\} \neq \mathfrak{P}_t(R)$ (i.e. R is not 1-Peirce), then there exists $\{e_1, 1 - e_1\} \subseteq \mathfrak{P}_t(R) \setminus \{0, 1\}$. By Theorem 4.2, $R^\pi \in \mathcal{T}_2$. Now if at least one of $e_1 R e_1$ or $(1 - e_1) R (1 - e_1)$ is not 1-Peirce, say $e_1 R e_1$, then there exists $e_2 \in \mathfrak{P}_t(e_1 R e_1) \setminus \{0, e_1\}$. By Lemma 3.8, $\{e_2, e_1 - e_2\} \subseteq \mathfrak{P}_t(e_1 R e_1) \subseteq \mathfrak{P}_{it}(e_1 R e_1) = e_1 R e_1 \cap \mathfrak{P}_{it}(R) \subseteq \mathfrak{P}_{it}(R)$. Hence $\{e_2, e_1 - e_2, 1 - e_1\}$ is a complete set of orthogonal idempotents contained in $\mathfrak{P}_{it}(R)$. By Theorem 4.2, $R^\pi \in \mathcal{T}_3$. If at least one of $e_2 R e_2$, $(e_1 - e_2) R (e_1 - e_2)$, or $(1 - e_1) R (1 - e_1)$ is not 1-Peirce, then either this inductive process will terminate in n steps for some $n \in \mathbb{Z}^+$ yielding condition (2) or it will continue indefinitely yielding condition (3). \square

Note that in Example 4.22(2), R has a 2-by-2 block form which is not in \mathcal{T}_2 . Observe that one can show that $E_{11} \in \mathfrak{P}_t(R)$, so R is not 1-Peirce. Surprisingly, Theorem 5.3 predicts that there is a 2-by-2 block form for R which is in \mathcal{T}_2 . Indeed, R can be partitioned into another 2-by-2 block form which is in \mathcal{T}_2 by taking R_1 to be a 1-by-1 matrix and R_2 to be a 3-by-3 matrix (i.e. this corresponds to taking $e_1 = E_{11}$ and $1 - e_1 = E_{22} + E_{33} + E_{44}$ in the proof of Theorem 5.3).

PROPOSITION 5.4 *Let $0 \neq e = e^2 \in R$ such that eRe is a 1-Peirce ring, $c \in \mathfrak{P}_t(R)$, $c_1 \in \mathfrak{P}_t(cRc)$, $0 \neq cec$ and $c_1 e c_1 \neq 0$. Then:*

- (1) $ece = e = ec_1e$.
- (2) $cecRcec$ is a 1-Peirce ring.

Proof (1) By Lemma 3.7, $0 \neq cec = (cec)^2 = c(ece)c$. Thus $ece \neq 0$. From Lemma 3.8(1), $c_1 \in \mathfrak{P}_{it}(R)$. So, by Lemma 3.7, $c_1 e c_1 = (c_1 e c_1)^2 = c_1 (e c_1 e) c_1$. Hence $ec_1e \neq 0$.

Claim 1. $ece \in \mathfrak{P}_{it}(eRe)$.

Let $x, y \in eRe$. By Lemma 3.7, $ece = (ece)^2$. Consider $(ece)x(e - ece)y(ece) = e(c[exe(1 - c)eye]c)e = e0e = 0$, because $c \in \mathfrak{P}_t(R) \subseteq \mathfrak{P}_{it}(R)$. Thus $ece \in \mathfrak{P}_{it}(eRe)$.

Claim 2. $ece \in \mathfrak{P}_t(eRe)$.

Consider $(e - ece)xecey(e - ece) = e[(1 - c)execeye(1 - c)]e = e0e = 0$, because $c \in \mathfrak{P}_{ot}(R)$. Thus $ece \in \mathfrak{P}_{ot}(eRe)$. Hence $0 \neq ece \in \mathfrak{P}_t(eRe)$. Since eRe is 1-Peirce, $ece = e$.

Claim 3. $ec_1e \in \mathfrak{P}_{it}(eRe)$.

From above, $c_1 \in \mathfrak{P}_{it}(R)$. Let $x, y \in eRe$. Then $ec_1ex(e - ec_1e)yec_1e = e[c_1xe(1 - c_1)eyc_1]e = 0$. Hence $ec_1e \in \mathfrak{P}_{it}(eRe)$.

Claim 4. $ec_1e \in \mathfrak{P}_t(eRe)$.

Since $e = ece$, we have

$$\begin{aligned} (e - ec_1e)xec_1ey(e - ec_1e) &= (ece - ec_1e)xc_1y(ece - ec_1e) \\ &= e(c - c_1)exc_1ye(c - c_1)e \\ &= e[(c - c_1)(cxc)c_1(cyc)(c - c_1)]e = 0, \end{aligned}$$

because $c_1 \in \mathfrak{P}_t(cRc) \subseteq \mathfrak{P}_{ot}(cRc)$. Thus $ec_1e \in \mathfrak{P}_t(eRe)$. Since eRe is a 1-Peirce ring, $ec_1e = e$.

(2) Let $0 \neq f \in \mathfrak{P}_t(cecRcec)$. Observe $0 \neq f = c[ecfce]c = c[efe]c$ since $c \in \mathfrak{P}_t(R)$. So $efe \neq 0$.

Claim 5. $efe = (efe)^2$.

Observe $f = (cec)f = c(cec)f = cf$. Similarly, $f = fc$. Consider $efe = e(fcec)f = efefe = (efe)^2$.

Claim 6. $efe \in \mathfrak{P}_{it}(eRe)$.

Let $x, y \in eRe$. By (1) $e = ece$, so

$$\begin{aligned} efex(e - efe)yefe &= efex(ece - efe)yefe = efexe(c - f)eyefe \\ &= e(fc)exe(c - f)eye(cf)e \\ &= ef[(cec)x(cec)(cec)(c - f)(cec)y(cec)]fe \\ &= ef[(cec)x(cec)f(cec)(c - f)(cec)y(cec)]fe \\ &= ef[(cec)x(cec)f(c - f)(cec)y(cec)]fe \\ &= ef[(cec)x(cec)0(cec)y(cec)]fe \\ &= 0, \end{aligned}$$

since $f \in \mathfrak{P}_t(cecRcec) \subseteq \mathfrak{P}_{it}(cecRcec)$.

Claim 7. $efe \in \mathfrak{P}_t(eRe)$.

Consider

$$\begin{aligned} (e - efe)xefey(e - efe) &= (ece - efe)xfy(ece - efe) \\ &= e(c - f)exfy(c - f)e \\ &= ece(c - f)xfy(c - f)ece \\ &= ecec(c - f)(cecxc)ef(cecyc)(c - f)ece \\ &= e(cec - f)[(cecxc)ef(cecyc)](cec - f)e \\ &= 0, \end{aligned}$$

since $f \in \mathfrak{P}_{ot}(cecRcec)$. Thus $efe \in \mathfrak{P}_t(eRe)$. Hence $efe = e$. So $0 \neq f = cecfcec = c(efe)c = cec$. Therefore $cecRcec$ is a 1-Peirce ring. \square

Observe that in Proposition 5.4 the conclusions $ece = e$ and $cecRcec$ is a 1-Peirce ring do not need the conditions $c_1 \in \mathfrak{P}_t(cRc)$ and $c_1ec_1 \neq 0$. Also, this result can be extended under related hypotheses (e.g. e primitive and $c \in \mathfrak{P}_{it}(R)$).

COROLLARY 5.5 *Let $0 \neq e = e^2 \in R$ such that eRe is a 1-Peirce ring and $c \in \mathfrak{P}_t(R)$. The following conditions are equivalent:*

- (1) $cec \neq 0$.

- (2) $ece = e$.
- (3) $(1 - c)e(1 - c) = 0$.

Proof

- (1) \Rightarrow (2) This implication follows from Proposition 5.4.
- (2) \Rightarrow (3) Since $c \in \mathfrak{P}_1(R)$, $e(1 - c)e = ece(1 - c)ece = 0$. By Lemma 3.7(1), $(1 - c)e(1 - c) = [(1 - c)e(1 - c)]^2 = (1 - c)e(1 - c)e(1 - c) = 0$.
- (3) \Rightarrow (1) Assume $(1 - c)e(1 - c) = 0$ and $cec = 0$. Then

$$\begin{aligned} e &= (c + (1 - c))e(c + (1 - c)) \\ &= cec + ce(1 - c) + (1 - c)ec + (1 - c)e(1 - c) \\ &= ce(1 - c) + (1 - c)ec. \end{aligned}$$

So $e = e^2 = (ce(1 - c) + (1 - c)ec)^2 = ce(1 - c)ce(1 - c) + ce(1 - c)ec + (1 - c)ece(1 - c) + (1 - c)ec(1 - c)ec = 0$, a contradiction. \square

COROLLARY 5.6 *Let $\{e_i\}_{i=1}^n$ be a complete set of nonzero orthogonal idempotents such that each $e_i Re_i$ is a 1-Peirce ring and $0 \neq c \in \mathfrak{P}_1(R)$. Then:*

- (1) $c = \sum_{i \in J_1} ce_i c$ and $1 - c = \sum_{i \in J_2} (1 - c)e_i(1 - c)$, where $ce_i c \neq 0$ for all $i \in J_1$ and $(1 - c)e_i(1 - c) \neq 0$ for all $i \in J_2$.
- (2) $|J_1| + |J_2| = n$.
- (3) $\{ce_i c \mid i \in J_1\} \cup \{(1 - c)e_i(1 - c) \mid i \in J_2\}$ is a complete set of orthogonal idempotents, where each $ce_i c R ce_i c$ and each $(1 - c)e_i(1 - c)R(1 - c)e_i(1 - c)$ is a 1-Peirce ring.

Proof

- (1) $c = c1c = c \sum_{i \in J_1} e_i c = \sum_{i \in J_1} ce_i c$, and similarly, $1 - c = \sum_{i \in J_2} (1 - c)e_i(1 - c)$, where $J_1 \cup J_2 \subseteq \{1, \dots, n\}$.
- (2) This part follows from Corollary 5.5.
- (3) Note that $1 = c + 1 - c = \sum_{i \in J_1} ce_i c + \sum_{i \in J_2} (1 - c)e_i(1 - c)$. Also, $(ce_i c)ce_j c = ce_i e_j c = 0$ for all $i \neq j$, since $c \in \mathfrak{P}_1(R)$. Similarly, $[(1 - c)e_i(1 - c)][(1 - c)e_j(1 - c)] = 0$ for all $i \neq j$. Moreover $[(1 - c)e_i(1 - c)][ce_j c] = 0 = [ce_i c][(1 - c)e_j(1 - c)]$ for all i, j . By Lemma 3.7, $ce_i c$ and $(1 - c)e_i(1 - c)$ are idempotents for all i . From Proposition 5.4(2), each $ce_i c R ce_i c$ and each $(1 - c)e_i(1 - c)R(1 - c)e_i(1 - c)$ is a 1-Peirce ring. \square

THEOREM 5.7

- (1) *If R is an n -Peirce ring ($n > 1$), then there is a complete set of orthogonal idempotents $\{e_i\}_{i=1}^n \subseteq \mathfrak{P}_{it}(R)$ (hence $R^\pi \in \mathcal{T}_n$) such that every $e_i Re_i$ is a 1-Peirce ring.*
- (2) *If a ring R has a complete set $\{e_i\}_{i=1}^n$ of orthogonal idempotents for some $n \geq 2$ such that every $e_i Re_i$ is a 1-Peirce ring, then R is a k -Peirce ring for some $1 \leq k \leq n$.*

Proof (1) We use strong induction on n . First, let R be a 2-Peirce ring. Then there is an $e \in \mathfrak{P}_t(R)$ such that eRe and $(1-e)R(1-e)$ are 1-Peirce rings, and $\{e, 1-e\}$ is a complete set of orthogonal idempotents, with $e, 1-e \in \mathfrak{P}_{it}(R)$.

Next, consider a fixed $n \geq 2$ and assume that for each $k, 2 \leq k \leq n$, if R is a k -Peirce ring, then there is a complete set of orthogonal idempotents $\{e_i\}_{i=1}^k \subseteq \mathfrak{P}_{it}(R)$ such that every e_iRe_i is a 1-Peirce ring. Now let R be an $(n+1)$ -Peirce ring. Then there is a $c \in \mathfrak{P}_t(R)$ such that cRc is a k -Peirce ring for some $k, 1 \leq k < n+1$, and $(1-c)R(1-c)$ is an $(n+1-k)$ -Peirce ring. Since $k \leq n$ and $n+1-k \leq n$, and assuming for the moment that $2 \leq k$ and $2 \leq n+1-k$, the induction hypothesis guarantees the existence of complete sets of orthogonal idempotents $\{e_i\}_{i=1}^k \subseteq \mathfrak{P}_{it}(cRc)$ and $\{f_j\}_{j=1}^{n+1-k} \subseteq \mathfrak{P}_{it}((1-c)R(1-c))$, such that $e_i c R c e_i$ and $f_j(1-c)R(1-c)f_j$ are 1-Peirce rings for every i and j . Since $c, 1-c \in \mathfrak{P}_{it}(R)$, it follows from Lemma 3.8 that $\{e_i\}_{i=1}^k, \{f_j\}_{j=1}^{n+1-k} \subseteq \mathfrak{P}_{it}(R)$. Since $(cRc)((1-c)R(1-c)) = 0$, we conclude that $\{e_i\}_{i=1}^k \cup \{f_j\}_{j=1}^{n+1-k}$ is an orthogonal set of idempotents in $\mathfrak{P}_{it}(R)$. Moreover, $\sum_{i=1}^k e_i + \sum_{j=1}^{n+1-k} f_j = c + (1-c) = 1$, and $e_i c R c e_i = e_i R e_i$ (since $e_i \in cRc$) and $f_j(1-c)R(1-c)f_j = f_j R f_j$ for every i and j .

Finally, we consider the case $k = 1$ or $n+1-k = 1$. Without loss of generality, let $k = 1$, i.e. $c \in \mathfrak{P}_{it}(R)$, cRc is a 1-Peirce ring and $(1-c)R(1-c)$ is an n -Peirce ring. Then we can proceed as in the previous paragraph with $c \in \mathfrak{P}_{it}(R)$ and a complete set of orthogonal idempotents $\{f_j\}_{j=1}^n \subseteq \mathfrak{P}_{it}((1-c)R(1-c))$, and then the set $\{c, f_1, \dots, f_n\}$ is a complete set of orthogonal idempotents in $\mathfrak{P}_{it}(R)$ such that cRc and $f_j R f_j$ are 1-Peirce rings for all j .

(2) We again use strong induction on n . Let R have a complete set of orthogonal idempotents $\{e_1, e_2\}$ such that each $e_i R e_i$ is a 1-Peirce ring. If R is a 1-Peirce ring, then we are done, since $1 \leq 2$. Otherwise there is a $c \in \mathfrak{P}_t(R)$ such that $c \notin \{0, 1\}$. Hence $1-c \in \mathfrak{P}_t(R)$ and $1-c \notin \{0, 1\}$. From Corollary 5.6, $c = c e_i c$ for $i \in \{1, 2\}$. Without loss of generality, assume $i = 1$. Then, again by Corollary 5.6, $cRc = c e_1 c R c e_1 c$, $(1-c)R(1-c) = (1-c)e_2(1-c)R(1-c)e_2(1-c)$, and cRc and $(1-c)R(1-c)$ are 1-Peirce rings. Therefore R is a 2-Peirce ring.

Next assume that the result holds for a fixed $n \geq 2$. Let R be a ring having a complete set of orthogonal idempotents $\{e_i\}_{i=1}^{n+1}$ such that each $e_i R e_i$ is a 1-Peirce ring. If R is a 1-Peirce ring, we are done. Otherwise there is a $c \in \mathfrak{P}_t(R)$ such that $c \notin \{0, 1\}$. Hence $1-c \in \mathfrak{P}_t(R)$ and $1-c \notin \{0, 1\}$. From Corollary 5.6, there exist $J_1, J_2 \subseteq \{1, \dots, n\}$ and complete sets of orthogonal idempotents $\{c e_i c \mid i \in J_1\}$ and $\{(1-c)e_i(1-c) \mid i \in J_2\}$ for cRc and $(1-c)R(1-c)$, respectively. From the induction hypothesis, there exist positive integers k_1 and k_2 such that $1 \leq k_1 \leq |J_1|$ and $1 \leq k_2 \leq |J_2|$ such that cRc is a k_1 -Peirce ring and $(1-c)R(1-c)$ is a k_2 -Peirce ring. Since $|J_1| + |J_2| = n+1$, then R is k -Peirce where $k = k_1 + k_2$ and $1 \leq k \leq n+1$. □

COROLLARY 5.8 *Let $\{e_i\}_{i=1}^n \subseteq \mathfrak{P}_{it}(R)$ be a complete set of orthogonal idempotents such that each $e_i R e_i$ is a k_i -Peirce ring for some positive integers k_i . Then R is a k -Peirce ring for some $1 \leq k \leq \sum_{i=1}^n k_i$.*

Proof This result follows from Theorem 5.7. □

From Theorem 5.7, R is an n -Peirce ($n > 1$) generalized matrix ring implies that $R \in \mathcal{T}_n$; and if R has a complete set of n orthogonal primitive idempotents, then R is k -Peirce for some k with $1 \leq k \leq n$. Thus it is natural to ask: If R has a complete set of orthogonal primitive idempotents $\{e_i\}_{i=1}^n \subseteq \mathfrak{P}_{it}(R)$ (hence $R^\pi \in \mathcal{T}_n$), must R be n -Peirce? Observe that for $n = 2$, the question has an affirmative answer. Our next example provides a negative answer, in general.

Example 5.9 Let

$$R = \begin{bmatrix} A & X^2 & X \\ X & A & X^2 \\ X^2 & X & A \end{bmatrix},$$

where A is a ring such that 0 and 1 are the only idempotents of A ; and $0 \neq X \triangleleft A$ such that $X \neq X^2$, $X^2 \neq X^3$ and $X^3 = 0$. Then $R \in \mathcal{T}_3$, but R is a 1-Peirce ring. One can construct such rings by letting B be a commutative ring, $0 \neq P$ a prime ideal of B such that $P \neq P^2$ and $P^2 \neq P^3$. Then take $A = B/P^3$ and $X = P/P^3$. In particular, let $B = F[x]$ where F is a field and $P = xF[x]$.

Since $X^2X = XX^2 = 0$, Corollary 4.3 yields $R \in \mathcal{T}_3$. To show that R is a 1-Peirce ring, we first characterize all nontrivial idempotents of R . Let $\alpha \in R$ such that $\alpha \neq 0$ and $\alpha \neq 1$. Then $\alpha = \alpha^2$ if and only if

$$\alpha = \begin{bmatrix} e_1 & m_{12} & m_{13} \\ m_{21} & e_2 & m_{23} \\ m_{31} & m_{32} & e_3 \end{bmatrix}$$

where $m_{12}, m_{23}, m_{31} \in X^2$, $m_{13}, m_{21}, m_{32} \in X$, $e_i \in \{0, 1\}$ and the following equations are satisfied:

$$\begin{aligned} e_1m_{12} + m_{12}e_2 + m_{13}m_{32} &= m_{12} \\ e_1m_{13} + m_{13}e_3 &= m_{13} \\ m_{21}e_1 + e_2m_{21} &= m_{21} \\ e_2m_{23} + m_{23}e_3 + m_{21}m_{13} &= m_{23} \\ m_{31}e_1 + e_3m_{31} + m_{32}m_{21} &= m_{31} \\ m_{32}e_2 + e_3m_{23} &= m_{32}. \end{aligned}$$

From the above conditions α must have one of the following six forms:

- (i) $\begin{bmatrix} 1 & m_{12} & m_{13} \\ m_{21} & 0 & m_{23} \\ m_{31} & 0 & 0 \end{bmatrix}$ with $m_{21}m_{13} = m_{23}$;
- (ii) $\begin{bmatrix} 0 & m_{12} & 0 \\ m_{21} & 1 & m_{23} \\ m_{31} & m_{32} & 0 \end{bmatrix}$ with $m_{32}m_{21} = m_{31}$;

- (iii) $\begin{bmatrix} 0 & m_{12} & m_{13} \\ 0 & 0 & m_{23} \\ m_{31} & m_{32} & 1 \end{bmatrix}$ with $m_{13}m_{32} = m_{12}$;
- (iv) $\begin{bmatrix} 1 & m_{12} & m_{13} \\ 0 & 1 & m_{23} \\ m_{31} & m_{32} & 0 \end{bmatrix}$ with $m_{13}m_{32} = -m_{12}$;
- (v) $\begin{bmatrix} 1 & m_{12} & 0 \\ m_{21} & 0 & m_{23} \\ m_{31} & m_{32} & 1 \end{bmatrix}$ with $m_{32}m_{21} = -m_{31}$;
- (vi) $\begin{bmatrix} 0 & m_{12} & m_{13} \\ m_{21} & 1 & m_{23} \\ m_{31} & 0 & 1 \end{bmatrix}$ with $m_{21}m_{13} = -m_{23}$.

Now assume R is not a 1-Peirce ring. Then there exist $c, 1 - c \in \mathfrak{P}_t(R)$ such that $0 \neq c$ and $c \neq 1$.

Then either c has a form of type (i), (ii) or (iii); or $1 - c$ has such a form. Without loss of generality, assume c has a form of type (i), (ii) or (iii). Then $1 - c$ has a form of type (iv), (v) or (vi). We show that no matrix of type (iv), (v) or (vi) is in $\mathfrak{P}_{ot}(R)$. Hence $1 - c \notin \mathfrak{P}_t(R)$, a contradiction. Therefore R is a 1-Peirce ring.

Since $X^2 \neq 0$, there exist $x, y \in X$ such that $0 \neq xy$.

Observe:

$$\begin{aligned} & \begin{bmatrix} 0 & m_{12} & m_{13} \\ m_{21} & 1 & m_{23} \\ m_{31} & 0 & 1 \end{bmatrix} xE_{21} \begin{bmatrix} 1 & -m_{12} & -m_{13} \\ -m_{21} & 0 & -m_{23} \\ -m_{31} & 0 & 0 \end{bmatrix} yE_{13} \begin{bmatrix} 0 & m_{12} & m_{13} \\ m_{21} & 1 & m_{23} \\ m_{31} & 0 & 1 \end{bmatrix} \\ &= xyE_{23} \neq 0; \\ & \begin{bmatrix} 1 & m_{12} & 0 \\ m_{21} & 0 & m_{23} \\ m_{31} & m_{32} & 1 \end{bmatrix} xE_{32} \begin{bmatrix} 0 & -m_{12} & 0 \\ -m_{21} & 1 & -m_{23} \\ -m_{31} & -m_{32} & 0 \end{bmatrix} yE_{21} \begin{bmatrix} 1 & m_{12} & 0 \\ m_{12} & 0 & m_{23} \\ m_{13} & m_{32} & 1 \end{bmatrix} \\ &= xyE_{31} \neq 0; \\ & \begin{bmatrix} 1 & m_{12} & m_{13} \\ 0 & 1 & m_{23} \\ m_{31} & m_{32} & 0 \end{bmatrix} xE_{13} \begin{bmatrix} 0 & -m_{12} & -m_{13} \\ 0 & 0 & -m_{23} \\ -m_{31} & -m_{32} & 1 \end{bmatrix} yE_{32} \begin{bmatrix} 1 & m_{12} & m_{13} \\ 0 & 1 & m_{23} \\ m_{31} & m_{32} & 0 \end{bmatrix} \\ &= xyE_{12} \neq 0. \end{aligned}$$

LEMMA 5.10 Let $0 \neq c = c^2 = R$ and $e \in \mathfrak{P}_{it}(cRc)$ such that $e \neq c$. Then $ReR \subsetneq RcR$.

Proof Observe that $\{e, c - e\}$ is set of orthogonal idempotents. Clearly, $ReR \subseteq RcR$. Assume that $ReR = RcR$. Then $c = \sum r_i e s_i$. So $c - e = (\sum r_i e s_i) - e$. Then

$$\begin{aligned} c - e &= (c - e)^2 = \left[\left(\sum r_i e s_i \right) - e \right] (c - e) \\ &= \left(\sum r_i e s_i \right) (c - e) \\ &= \sum r_i e s_i (c - e) \end{aligned}$$

$$\begin{aligned}
 &= \left(\sum r_i e s_i (c - e) \right)^2 \\
 &= \sum_j \sum_i r_i e s_i (c - e) r_j e s_j (c - e) \\
 &= 0,
 \end{aligned}$$

since

$$\begin{aligned}
 r_i [e s_i (c - e) r_j e] s_j (c - e) &= r_i [e s_i e (c - e) r_j e] s_j (c - e) \\
 &= r_i 0 s_j (c - e) \\
 &= 0,
 \end{aligned}$$

because $e \in \mathfrak{P}_{it}(cRc)$. However, this is a contradiction, since $e \neq c$. Therefore $ReR \subsetneq RcR$. □

THEOREM 5.11 *Assume R has DCC on $\{ReR \mid e \in \mathfrak{P}_t(R)\}$. Then R is an n -Peirce ring for some $n \in \mathbb{Z}^+$.*

Proof Assume R has DCC on $\{ReR \mid e \in \mathfrak{P}_t(R)\}$, but R is not an n -Peirce ring for any $n \in \mathbb{Z}^+$. Observe that $\mathfrak{P}_t(R) \neq \{0, 1\}$. So let $0 \neq c_1 \in \mathfrak{P}_t(R)$ be such that $c_1 \neq 1$. Then $c_1 R c_1$ is not an n -Peirce ring for any $n \in \mathbb{Z}^+$, or $(1 - c_1)R(1 - c_1)$ is not an n -Peirce ring for any $n \in \mathbb{Z}^+$. Without loss of generality, say $c_1 R c_1$ is not an n -Peirce ring for any $n \in \mathbb{Z}^+$. Then $\mathfrak{P}_t(c_1 R c_1) \neq \{0, c_1\}$. So let $0 \neq c_2 \in \mathfrak{P}_t(c_1 R c_1)$ be such that $c_2 \neq c_1$. Then $c_2 R c_2$ is not an n -Peirce ring for any $n \in \mathbb{Z}^+$, or $(c_1 - c_2)R(c_1 - c_2)$ is not an n -Peirce ring for any $n \in \mathbb{Z}^+$. Without loss of generality, say $c_2 R c_2$ is not an n -Peirce ring for any $n \in \mathbb{Z}^+$. By Lemma 3.8, $c_1, c_2 \in \mathfrak{P}_{it}(R)$. From Lemma 5.10, $R \supsetneq Rc_1R \subsetneq Rc_2R$. We can continue this process indefinitely, which contradicts the DCC on $\{ReR \mid e \in \mathfrak{P}_{it}(R)\}$. Therefore R is an n -Peirce ring for some $n \in \mathbb{Z}^+$. □

PROPOSITION 5.12 *If $\{b_1, \dots, b_n\} \subseteq \mathfrak{P}_{it}(R)$ is a complete set of nonzero orthogonal idempotents such that $\mathfrak{P}_{it}(b_i R b_i) = \{0, b_i\}$, then $|\{ReR \mid e \in \mathfrak{P}_{it}(R)\}| \leq 2^n$.*

Proof Let $0 \neq e = e^2 \in \mathfrak{P}_{it}(R)$. By Lemma 3.8, each $b_i e b_i \in \mathfrak{P}_{it}(R)$. Observe that $e = (\sum_{i=1}^n b_i) e (\sum_{i=1}^n b_i) = \sum_{i=1}^n \sum_{j=1}^n b_i e b_j$, and $b_i e b_i \neq 0$ if and only if $b_i e b_i = b_i$. Let $J \subseteq \{1, \dots, n\}$ such that $b_i e b_i \neq 0$ if and only if $i \in J$. Since $e \in \mathfrak{P}_{it}(R)$, $e = e(e)e = e(\sum_{i=1}^n \sum_{j=1}^n b_i e b_j)e = \sum_{i=1}^n \sum_{j=1}^n e b_i e b_j e = \sum_{i=1}^n \sum_{j=1}^n e b_i b_j e = \sum_{i \in J} e b_i e$. Then $ReR \subseteq \sum_{i \in J} R b_i R$.

Observe that $e - \sum_{i \neq j} b_i e b_j = \sum_{i \in J} b_i e b_i = \sum_{i \in J} b_i$. Let $k \in J$, then $b_k e - b_k e b_j = b_k$. Hence $R b_k R \subseteq ReR$. So $\sum_{i \in J} R b_i R \subseteq ReR$. Therefore $ReR = \sum_{i \in J} R b_i R$. Since $J \subseteq \{1, \dots, n\}$, $|\{ReR \mid e \in \mathfrak{P}_{it}(R)\}| \leq |\{\sum_{i \in K} R b_i R \mid K \subseteq \{1, \dots, n\}\}| = |\{K \mid K \subseteq \{1, \dots, n\}\}| = 2^n$, where $\sum_{i \in K} R b_i R$ corresponds to $\{0\}$ when $K = \emptyset$. □

As an illustration and application of several of our previous results, we provide the following lemma and proposition. Recall that a ring R is ‘quasi-Baer’ if for each $X \trianglelefteq R$ there is an $e = e^2 \in R$ such that $r_R(X) = eR$. See [6,21] for further details on the class of quasi-Baer rings.

LEMMA 5.13 *R is a prime ring if and only if R is quasi-Baer and a 1-Peirce ring.*

Proof Assume R is prime. From [3, Lemma 4.2] or [6, Proposition 3.2.5], R is quasi-Baer. From Corollary 3.6, R is a 1-Peirce ring. Conversely, assume $xRy = 0$ for some $x, y \in R$ with $x \neq 0$. Then $y \in \underline{r}_R(xR) = \underline{r}_R(RxR) = eR$ for some $e = e^2 \in R$. Since $\underline{r}_R(xR)$ is an ideal of R , then $e \in \mathcal{S}_\ell(R)$. By Proposition 3.10(1), $e = 0$. Hence $y = 0$, so R is prime. \square

PROPOSITION 5.14 *Assume that R is a quasi-Baer ring. If $\{e_1, \dots, e_n\}$ is a complete set of orthogonal inner Peirce trivial idempotents and each $e_i R e_i$ is a 1-Peirce ring, then R is a k-Peirce ring for some $1 \leq k \leq n$, $R^\pi \in \mathcal{T}_n$ and each $e_i R e_i$ is a prime ring.*

Proof This proof follows from Theorem 5.7(2), Theorem 4.2, Lemma 5.13, and [21, Lemma 2]. \square

For example, any quasi-Baer ring with a complete set of orthogonal primitive idempotents (e.g. a right hereditary right Noetherian ring) satisfies the hypothesis of Proposition 5.14.

In a sequel to this paper, we further investigate the properties and structure of the class of n -Peirce rings.

Acknowledgements

Proposition 4.8 was suggested by Arturo Magidin. Our attention was drawn to [2] by Lance Small. We are grateful to the referee for his/her efforts.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

The first author was supported by the Hungarian National Foundation for Scientific Research under [grants number K-101515] and Institute of Mathematics, Hanoi, Vietnam. The third author was supported by the National Research Foundation (of South Africa) under [grant number UID 72375]. Any opinion, findings and conclusions or recommendations expressed in this material are those of the authors and therefore the National Research Foundation does not accept any liability in regard thereto.

References

- [1] Krylov PA, Tuganbaev AA. Modules over formal matrix rings. *Fundam. Prikl. Mat.* 2009;15(8):145–211. translation in *J. Math. Sci. (NY)*. (2010);171(2):248–295. Russian.
- [2] Peirce B. Linear associative algebra. *Am. J. Math.* 1881;4:97–229.
- [3] Birkenmeier GF, Heatherly HE, Kim JY, Park JK. Triangular matrix representations. *J. Algebra.* 2000;230:558–595.
- [4] Ánh PN, van Wyk L. Automorphism groups of generalized triangular matrix rings. *Linear Algebra Appl.* 2011;434:1018–1025.

- [5] Ánh PN, van Wyk L. Isomorphisms between strongly triangular matrix rings. *Linear Algebra Appl.* 2013;438:4374–4381.
- [6] Birkenmeier GF, Park JK, Rizvi ST. *Extensions of rings and modules*. New York (NY): Birkhäuser/Springer; 2013.
- [7] Rowen LH. *Ring theory I*. Boston (MA): Academic Press; 1988.
- [8] Birkenmeier GF, Park JK, Rizvi ST. Hulls of semiprime rings with applications to C^* -algebras. *J. Algebra.* 2009;322:327–352.
- [9] Fisher JW, Snider RL. On the Von Neumann regularity of rings with regular prime factor rings. *Pac. J. Math.* 1974;54:135–144.
- [10] Gardner BJ, Wiegandt R. *Radical theory of rings*. New York (NY): Marcel Dekker; 2004.
- [11] Tang G, Li C, Zhou Y. Study of Morita contexts. *Commun. Algebra.* 2014;42:1668–1681.
- [12] Szigeti J. Linear algebra in lattices and nilpotent endomorphisms of semisimple modules. *J. Algebra.* 2008;319:296–308.
- [13] Armendariz EP, Birkenmeier GF, Park JK. Ideal intrinsic extensions with connections to PI-rings. *J. Pure Appl. Algebra.* 2009;213:1756–1776; Corrigendum. 2011;215:99–100.
- [14] Wisbauer R. *Modules and algebras: bimodule structure and group actions on algebras*. Harlow: Addison Wesley Longman; 1996.
- [15] Rowen LH. Examples of semiperfect rings. *Isr. J. Math.* 1989;65:273–283.
- [16] Cedo F, Rowen LH. Addendum to “Examples of semiperfect rings”. *Isr. J. Math.* 1998;107:343–348.
- [17] Dung NV, Huynh DV, Smith PF, Wisbauer R. *Extending modules*. Harlow: Longman; 1994.
- [18] Birkenmeier GF, Müller BJ, Rizvi ST. Modules in which every fully invariant submodule is essential in a direct summand. *Commun. Algebra.* 2002;30:1395–1415.
- [19] Kaplansky I. *Rings of operators*. New York (NY): Benjamin; 1968.
- [20] Lam TY. *Lectures on modules and rings*. New York (NY): Springer; 1999.
- [21] Clark WE. Twisted matrix units semigroup algebras. *Duke Math. J.* 1967;34:417–423.