

The Fibonacci number of Fibonacci trees and a related family of polynomial recurrence systems.

Stephan G. Wagner

Department of Mathematics

Graz University of Technology

Steyrergasse 30, A-8010 Graz, Austria

wagner@finanz.math.tu-graz.ac.at

Abstract

Fibonacci trees are special binary trees which are of natural interest in the study of data structures. A Fibonacci tree of order n has the Fibonacci trees of orders $n - 1$ and $n - 2$ as left and right subtrees. On the other hand, the Fibonacci number $F(G)$ of a graph G , introduced in a paper of Prodinger and Tichy in 1982, is defined as the number of independent vertex subsets of G . In this paper, we study the Fibonacci number of Fibonacci trees and show that the underlying system of recurrence equations belongs to a class with a special property. It will be shown that the Fibonacci number of the n -th Fibonacci tree with $F_{n+2} - 1$ vertices is asymptotically $0.682328 \cdot (3.659873)^{F_n}$.

1 Introduction and Preliminaries

Fibonacci trees (cf. [4, 7]) are special balanced trees which are of natural interest in computer science. They are defined as binary trees, where the

Fibonacci tree of order n has a left subtree which is a Fibonacci tree of order $n - 1$ and a right subtree of order $n - 2$ (an order 0 tree has no nodes, an order 1 tree has exactly 1 node). The Fibonacci tree of order n has exactly $F_{n+2} - 1$ nodes.

Fibonacci trees appear in the study of AVL-trees, special highly balanced binary trees. The heights of the two subtrees at any vertex of an AVL-tree may only differ by at most 1. A Fibonacci tree is thus the most unbalanced AVL-tree that is allowed. Figure 1 shows the Fibonacci tree of order 4.




Figure 1: The Fibonacci tree of order 4.

On the other hand, the concept of the Fibonacci number for graphs was introduced in a paper of Prodinger and Tichy [9]. The Fibonacci number $F(G)$ of a graph G is defined as the number of independent vertex subsets of G , where a set of vertices is said to be independent if it contains no pair of connected vertices. The name is due to the fact that $F(P_n)$, where P_n is a simple path of length n , gives the sequence of Fibonacci numbers. Similarly, $F(C_n)$, where C_n denotes the cycle of length n , gives the Lucas numbers.

In the original work, the authors could prove that the star has maximal Fibonacci number among all trees, whereas the path has minimal Fibonacci number. They also considered recursions for several classes of graphs. In subsequent papers ([5, 6]), classes of simply generated trees were investigated. The concept of Fibonacci numbers even turned out to be of use in combinatorial chemistry (cf. [8]).

For a true Fibonacci enthusiast, it is very natural to bring the two concepts

together by considering the Fibonacci number of a Fibonacci tree. For this purpose, we deduce a system of recurrence equations. Let a_n be the number of independent subsets of the n -th Fibonacci tree which contain the root and let b_n be the number of independent subsets which do not contain the root. It is quite obvious now – from the construction of the Fibonacci trees – that the equations

$$a_n = b_{n-1}b_{n-2}$$

and

$$b_n = (a_{n-1} + b_{n-1})(a_{n-2} + b_{n-2})$$

hold. Together with the initial values $a_0 = 0$, $a_1 = b_0 = b_1 = 1$, these equations determine a_n and b_n uniquely. Recurrences of this type typically lead to a doubly exponential growth (cf. [1]), and this also holds in our case. However, it will be shown in the following section that the fraction $\frac{a_n}{b_n}$ can be given by an explicit formula in terms of n , which is quite unusual for recurrences of this kind; this phenomenon is due to the special form of the recurrence; we will generalize the recurrence to a parametric class for which this property still holds. Moreover, we will derive asymptotic expressions for a_n , b_n and the total number of independent vertices, $a_n + b_n$.

2 A class of recurrence equations and its properties

We generalize the system for the Fibonacci number of Fibonacci trees in the following way: let α and β be arbitrary positive numbers, and fix an integer $k > 1$. Now, let us consider the system of recurrences given by

$$\begin{aligned} a_n &= \prod_{i=1}^k b_{n-i}, \\ b_n &= \prod_{i=1}^k (\alpha a_{n-i} + \beta b_{n-i}), \end{aligned} \tag{1}$$

together with positive initial values a_i, b_i ($0 \leq i < k$). First, we are going to prove a short lemma on the quotient $\frac{a_n}{b_n}$ – quite surprisingly, we can give a rather simple formula for this fraction in terms of n :

Lemma 1 Let a_n, b_n be given by the system (1). Then, the quotient $\frac{a_n}{b_n}$ can be written as $\frac{a_n}{b_n} = \frac{r_n}{r_{n+k}}$, where r_n is a linear recurrent sequence given by the initial values $r_n = \frac{a_n}{b_n} \prod_{j=0}^{n-1} \left(\frac{\alpha a_j}{b_j} + \beta \right)$ ($0 \leq n \leq k$) and the recurrence $r_n = \beta r_{n-1} + \alpha r_{n-k-1}$.

Proof: By simple induction on n . Note first that

$$\frac{b_k}{a_k} = \prod_{j=0}^{k-1} \frac{\alpha a_j + \beta b_j}{b_j} = \prod_{j=0}^{k-1} \left(\frac{\alpha a_j}{b_j} + \beta \right)$$

by (1), so $r_k = 1$. Thus, we have, for $n \leq k$,

$$\begin{aligned} r_{n+k} &= \beta r_{n+k-1} + \alpha r_{n-1} = \beta^2 r_{n+k-2} + \alpha r_{n-1} + \alpha \beta r_{n-2} \\ &= \dots = \beta^n r_k + \alpha \sum_{j=1}^n \beta^{j-1} r_{n-j} \\ &= \beta^n + \alpha \sum_{j=1}^n \beta^{j-1} \frac{a_{n-j}}{b_{n-j}} \prod_{i=0}^{n-j-1} \left(\frac{\alpha a_i}{b_i} + \beta \right), \end{aligned}$$

and it is not difficult to prove (by means of another induction) that this equals

$$\prod_{j=0}^{n-1} \left(\frac{\alpha a_j}{b_j} + \beta \right).$$

Therefore,

$$\frac{r_n}{r_{n+k}} = \frac{a_n}{b_n}$$

holds for $n \leq k$. For the induction step, note again that dividing the two equations in (1) yields

$$\frac{b_n}{a_n} = \prod_{i=1}^k \left(\alpha \frac{a_{n-i}}{b_{n-i}} + \beta \right).$$

Now, by the definition of r_n and the induction hypothesis, we have

$$\alpha \frac{a_{n-i}}{b_{n-i}} + \beta = \frac{\alpha r_{n-i} + \beta r_{n-i+k}}{r_{n-i+k}} = \frac{r_{n-i+k+1}}{r_{n-i+k}}$$

and hence

$$\prod_{i=1}^k \left(\alpha \frac{a_{n-i}}{b_{n-i}} + \beta \right) = \left(\prod_{i=n+1}^{n+k} r_i \right) \left(\prod_{i=n}^{n+k-1} r_i \right)^{-1} = \frac{r_{n+k}}{r_n},$$

which finishes the induction. \square

Now, the first equation in (1) can be written as

$$x_n = \sum_{i=1}^k x_{n-i} + y_n,$$

where $x_n = \log b_n$ and $y_n = \log \frac{b_n}{a_n} = \log \frac{r_{n+k}}{r_n}$. Iteration yields

$$x_n = \sum_{i=k}^n d_{k-1, n+k-i-1} y_i + \sum_{l=0}^{k-1} d_{l,n} x_l,$$

where the sequences $d_{l,n}$ are defined by $d_{l,n} = 0$ ($0 \leq n < k$, $n \neq l$), $d_{l,l} = 1$ and $d_{l,n} = \sum_{i=1}^k d_{l, n-i}$. Note that we obtain the Fibonacci sequence in the case $k = 2$. From a result of Brauer [2], we know that the polynomial $x^k - \sum_{i=0}^{k-1} x^i$ has a dominant root $\phi > 1$ and all other roots have absolute value < 1 . Let $\phi = \phi_1, \phi_2, \phi_3, \dots, \phi_k$ be the roots of this polynomial. Then there are constants $\gamma_{l,j}$ ($0 \leq l < k$, $1 \leq j \leq k$) such that

$$d_{l,n} = \sum_{j=1}^k \gamma_{l,j} \phi_j^n.$$

Suppose that ϕ_2 is the root of second-largest modulus. Then

$$d_{l,n} = \gamma_{l,1} \phi^n + O(|\phi_2|^n)$$

and thus

$$\sum_{l=0}^{k-1} d_{l,n} x_l = \left(\sum_{l=0}^{k-1} \gamma_{l,1} x_l \right) \phi^n + O(|\phi_2|^n).$$

Furthermore,

$$\sum_{i=k}^n d_{k-1, n+k-i-1} y_i = \sum_{j=1}^k \gamma_{k-1,j} \sum_{i=k}^n \phi_j^{n+k-i-1} y_i.$$

We have to distinguish two different cases for the inner sum. For both cases, we note first that the polynomial $x^{k+1} - \beta x^k - \alpha$ has a dominant positive root

ψ , so $r_n = \delta\psi^n(1 + O(\eta^{-n}))$, where δ and $\eta > 1$ are positive constants. This shows that $y_n = \log \frac{r_{n+k}}{r_n} = (k \log \psi)(1 + O(\eta^{-n}))$. We use C as an abbreviation for $k \log \psi$. Now, for $j > 1$, we have

$$\begin{aligned}
\sum_{i=k}^n \phi_j^{n+k-i-1} y_i &= \sum_{i=k-1}^{n-1} \phi_j^i y_{n+k-1-i} \\
&= \sum_{k-1 \leq i \leq \frac{n+k}{2}-1} \phi_j^i y_{n+k-1-i} + \sum_{\frac{n+k}{2}-1 < i \leq n-1} \phi_j^i y_{n+k-1-i} \\
&= \sum_{k-1 \leq i \leq \frac{n+k}{2}-1} \phi_j^i C(1 + O(\eta^{-n/2})) + O\left(\sum_{\frac{n+k}{2}-1 < i \leq n-1} |\phi_j|^i\right) \\
&= C \sum_{i=k-1}^{\infty} \phi_j^i + O(|\phi_j|^{n/2}) + O(\eta^{-n/2}).
\end{aligned}$$

On the other hand,

$$\begin{aligned}
\sum_{i=k}^n \phi^{n+k-i-1} y_i &= \phi^{n+k-1} \sum_{i=k}^n \phi^{-i} y_i \\
&= \phi^{n+k-1} \sum_{i=k}^{\infty} \phi^{-i} y_i - \phi^{n+k-1} \sum_{i=n+1}^{\infty} \phi^{-i} y_i \\
&= \phi^{n+k-1} \sum_{i=k}^{\infty} \phi^{-i} y_i - C \phi^{n+k-1} \sum_{i=n+1}^{\infty} \phi^{-i} (1 + O(\eta^{-n})) \\
&= \phi^{n+k-1} \sum_{i=k}^{\infty} \phi^{-i} y_i - \frac{C \phi^{k-1}}{\phi - 1} + O(\eta^{-n}).
\end{aligned}$$

Altogether, we see that there are positive constants A, B such that

$$b_n \sim A \cdot B^{\phi^n} \text{ and } a_n \sim e^{-C} A \cdot B^{\phi^n}.$$

Inserting in the first equation of (1) yields $A = e^{-\frac{C}{k-1}}$, so we arrive at the following theorem:

Theorem 2 *If a_n, b_n are given by the system (1), then there is a positive constant B (which depends on the initial values) and another constant $\theta = \min(|\phi_2|^{-1/2}, \eta^{1/2}) > 1$ such that*

$$b_n = \psi^{-\frac{k}{k-1}} \cdot B^{\phi^n} (1 + O(\theta^{-n})) \text{ and } a_n = \psi^{-\frac{k^2}{k-1}} \cdot B^{\phi^n} (1 + O(\theta^{-n})),$$

where ψ is the unique positive root of the polynomial $x^{k+1} - \beta x^k - \alpha$ and ϕ is the dominant root of $x^k - \sum_{i=0}^{k-1} x^i$.

REMARK: In the special case that corresponds to the Fibonacci numbers of Fibonacci trees (i.e. $k = 2$, $\alpha = \beta = 1$), we have $\psi = 1.465571\dots$ and $B = 1.786445\dots$. B is easily calculated by the formula

$$\log B = \frac{1}{\sqrt{5}} \sum_{i=0}^{\infty} \left(\frac{1 + \sqrt{5}}{2} \right)^{-i} \log \frac{r_{i+2}}{r_i} = \frac{1}{\sqrt{5}} \sum_{i=1}^{\infty} \left(\frac{1 + \sqrt{5}}{2} \right)^{-i} r_{i+1},$$

where $r_0 = r_1 = r_2 = 1$ and $r_n = r_{n-1} + r_{n-3}$. Now, we have

$$a_n \sim A_1 B^{\phi^n}, \quad b_n \sim A_2 B^{\phi^n},$$

where A_1 and A_2 are the positive roots of the polynomials $x^3 - 2x^2 + 5x - 1$ and $x^3 + 2x^2 + x - 1$ respectively. If we set $B' := B^{\sqrt{5}}$, we obtain

$$a_n \sim A_1 B'^{F_n}, \quad b_n \sim A_2 B'^{F_n},$$

where F_n denotes the Fibonacci numbers. Thus, a_n and b_n almost behave like the “multiplicative Fibonacci sequences” studied in [3]. Remembering that the number of vertices of the n -th Fibonacci tree is exactly $V_n = F_{n+2} - 1$, we see that the Fibonacci number of the n -th Fibonacci tree is asymptotically

$$1.120000 \cdot (1.641439)^{V_n}.$$

From [9], we know that the Fibonacci number of a tree with V vertices lies between F_{V+2} and $2^{V-1} + 1$. Furthermore, it was shown in [10] that the average Fibonacci number of a tree is asymptotically $1.129410 \cdot (1.674490)^V$, so the Fibonacci number of a Fibonacci tree is comparatively small.

REMARK: We can easily modify the construction of Fibonacci trees in the following way: the subtrees of the k -ary analogue of order n are the trees of order $n - 1, n - 2, \dots, n - k$. Then, the corresponding system of recurrences for the Fibonacci numbers of these trees has exactly the form (1).

Acknowledgement

The author was supported by Austrian Science Fund project no. S-8307-MAT.

References

- [1] A. V. Aho and N. J. A. Sloane. Some doubly exponential sequences. *Fibonacci Quart.*, 11(4):429–437, 1973.
- [2] A. Brauer. On algebraic equations with all but one root in the interior of the unit circle. *Math. Nachr.*, 4:250–257, 1951.
- [3] S. Falcon. Fibonacci’s multiplicative sequence. *Internat. J. Math. Ed. Sci. Tech.*, 34(2):310–315, 2003.
- [4] R. P. Grimaldi. Properties of Fibonacci trees. In *Proceedings of the Twenty-second Southeastern Conference on Combinatorics, Graph Theory, and Computing (Baton Rouge, LA, 1991)*, volume 84, pages 21–32, 1991.
- [5] P. Kirschenhofer, H. Prodinger, and R. F. Tichy. Fibonacci numbers of graphs. II. *Fibonacci Quart.*, 21(3):219–229, 1983.
- [6] P. Kirschenhofer, H. Prodinger, and R. F. Tichy. Fibonacci numbers of graphs. III. Planted plane trees. In *Fibonacci numbers and their applications (Patras, 1984)*, volume 28 of *Math. Appl.*, pages 105–120. Reidel, Dordrecht, 1986.
- [7] D. E. Knuth. *The art of computer programming. Volume 3*. Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1973. Sorting and searching, Addison-Wesley Series in Computer Science and Information Processing.
- [8] R. E. Merrifield and H. E. Simmons. *Topological Methods in Chemistry*. Wiley, New York, 1989.
- [9] H. Prodinger and R. F. Tichy. Fibonacci numbers of graphs. *Fibonacci Quart.*, 20(1):16–21, 1982.

- [10] S. G. Wagner. Subset counting in trees. *Ars Combinatoria*, to appear, 2006.

AMS Classification Numbers: 05C30, 05C69, 11B37